

<https://helda.helsinki.fi>

Fundamental research questions in subterranean biology

Mammola, Stefano

2020-12

Mammola , S , Amorim , I R , Bichuette , M E , Borges , P A V , Cheeptham , N , Cooper , S J B , Culver , D C , Deharveng , L , Eme , D , Ferreira , R L , Fiser , C , Fiser , Z , Fong , D W , Griebler , C , Jeffery , W R , Jugovic , J , Kowalko , J E , Lilley , T M , Malard , F , Manenti , R , Martinez , A , Meierhofer , M B , Niemiller , M L , Northup , D E , Pellegrini , T G , Pipan , T , Protas , M , Reboleira , A S P S , Venarsky , M P , Wynne , J J , Zagmajster , M & Cardoso , P 2020 , ' Fundamental research questions in subterranean biology ' , Biological Reviews , vol. 95 , no. 6 , pp. 1855-1872 . <https://doi.org/10.1111/brv.12642>

<http://hdl.handle.net/10138/333518>

<https://doi.org/10.1111/brv.12642>

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

This is the accepted version of the following article: FULL CITE, which has been published in final form at [10.1111/brv.12642]. This article may be used for non-commercial purposes in accordance with the Wiley Self-Archiving Policy [http://www.wileyauthors.com/self-archiving].

Fundamental research questions in subterranean biology

Stefano Mammola^{1,2,*}, Isabel R. Amorim³, Maria E. Bichuette⁴, Paulo A. V. Borges³, Naowarat Cheeptham⁵, Steven J.B. Cooper^{6,7}, David C. Culver⁸, Louis Deharveng⁹, David Eme¹⁰, Rodrigo Lopes Ferreira¹¹, Cene Fišer¹², Žiga Fišer¹², Daniel W. Fong¹³, Christian Griebler¹⁴, William R. Jeffery¹⁵, Jure Jugovic¹⁶, Johanna E. Kowalko¹⁷, Thomas M Lilley¹⁸, Florian Malard¹⁹, Raoul Manenti²⁰, Alejandro Martínez², Melissa B. Meierhofer^{18,21}, Matthew L. Niemiller²², Diana E. Northup²³, Thais G. Pellegrini¹¹, Tanja Pipan^{24,25}, Meredith Protas²⁶, Ana Sofia P. S. Reboleira²⁷, Michael P. Venarsky²⁸, J. Judson Wynne²⁹, Maja Zagamajster¹² and Pedro Cardoso^{1,*}

¹Laboratory for Integrative Biodiversity Research (LIBRe), Finnish Museum of Natural History (LUOMUS), University of Helsinki, Pohjoinen Rautatiekatu 13, 00100 Helsinki, Finland

²Molecular Ecology Group (MEG), Water Research Institute (IRSA), National Research Council (CNR), Corso Tonolli, 50, 28922 Pallanza, Italy

22 ³cE3c – Centre for Ecology, Evolution and Environmental Changes / Azorean Biodiversity
 23 Group and Universidade dos Açores, Faculty of Agrarian and Environmental Sciences. Rua
 24 Capitão João d'Àvila, Pico da Urze, 9700-042 Angra do Heroísmo, Azores, Portugal.
 25 ⁴Laboratory of Subterranean Studies, Federal University of São Carlos, Rodovia Washington
 26 Luís km 235, 13565-905 São Carlos, São Paulo, Brazil
 27 ⁵Department of Biological Sciences, Faculty of Science, Thompson Rivers University, 805 TRU
 28 Way, Kamloops, BC, Canada
 29 ⁶Evolutionary Biology Unit, South Australian Museum, North Terrace, Adelaide, South Australia
 30 5000, Australia
 31 ⁷Australian Centre for Evolutionary Biology and Biodiversity, and Environment Institute, School
 32 of Biological Sciences, University of Adelaide, Adelaide, South Australia 5005, Australia
 33 ⁸Department of Environmental Science, American University, 4400 Massachusetts Avenue, N.W.,
 34 Washington DC 20016, USA
 35 ⁹UMR7205 – ISYEB, Museum national d'Histoire naturelle, 45 rue Buffon (CP50), 75005 Paris,
 36 France
 37 ¹⁰IFREMER centre Atlantique, unité Ecologie et Modèles pour l'Halieutique, Rue de l'Île d'Yeu,
 38 44980, Nantes, France
 39 ¹¹Center of Studies in Subterranean Biology, Biology Department, Federal University of Lavras,
 40 Campus Universitário, CEP 37202-553, Lavras, Minas Gerais, Brazil
 41 ¹²SubBio Lab, Department of Biology, Biotechnical Faculty, University of Ljubljana,
 42 Jamnikarjeva 101, Ljubljana, SI-1000, PO BOX 2995, Slovenia
 43 ¹³Department of Biology, American University, 4400 Massachusetts Avenue, N.W.
 44 Washington DC 20016, USA

45 ¹⁴*University of Vienna, Department of Functional and Evolutionary Ecology, Division of*
46 *Limnology, Althanstrasse 14, 1090 Vienna*

47 ¹⁵*Department of Biology, University of Maryland, College Park, MD, Maryland 20742, USA*

48 ¹⁶*Department of Biodiversity, Faculty of Mathematics, Natural Sciences and Information*
49 *Technologies, University of Primorska, Glagoljaška 8, Koper, SI-6000, Slovenia*

50 ¹⁷*Harriet L. Wilkes Honors College, Florida Atlantic University, 5353 Parkside Dr, Jupiter, FL*
51 *33458, USA*

52 ¹⁸*BatLab Finland, Finnish Museum of Natural History, University of Helsinki, Pohjoinen*
53 *Rautatiekatu 13, 00100 Helsinki, Finland*

54 ¹⁹*UMR5023 Ecologie des Hydrosystèmes Naturels et Anthropisés, Univ. Lyon 1, ENTPE, CNRS,*
55 *Univ. de Lyon, Bat. Forel, 6 rue Raphaël Dubois, 69622 Villeurbanne cedex, France*

56 ²⁰*Department of Environmental Science and Policy, Università degli Studi di Milano, Via*
57 *Celoria 26, 20113 Milano, Italy*

58 ²¹*Department of Rangeland, Wildlife and Fisheries Management, Texas A&M University, 534*
59 *John Kimbrough Blvd., College Station, TX 77843, USA*

60 ²²*Department of Biological Sciences, The University of Alabama in Huntsville, 301 Sparkman*
61 *Drive NW, Huntsville, AL 35899, USA*

62 ²³*Department of Biology, University of New Mexico, Albuquerque, NM 87131-0001, USA*

63 ²⁴*ZRC SAZU Karst Research Institute, Novi trg 2, SI-1000 Ljubljana, Slovenia*

64 ²⁵*UNESCO Chair on Karst Education, University of Nova Gorica, Vipavska cesta, 5000 Nova*
65 *Gorica, Slovenia*

66 ²⁶*Department of Natural Sciences and Mathematics, Domenicas University of California, 50*
67 *Acacia Avenue, San Rafael, CA 94901, USA*

²⁷*Natural History Museum of Denmark, University of Copenhagen, Universitetsparken 15, 2100 Copenhagen, Denmark*

²⁸*Australian Rivers Institute, Griffith University, 170 Kessels Road, Nathan 4111, Queensland, Australia*

²⁹*Department of Biological Sciences, Center for Adaptable Western Landscapes, Northern Arizona University, Box 5640, Flagstaff, AZ 86011, US*

Running title: Scanning the horizon of subterranean biology

*Authors for correspondence: S. Mammola (E-mail: stefano.mammola@helsinki.fi; stefano.mammola@cnr.it; Tel.: +39 0323 518363; Water Research Institute, National Research Council of Italy, Corso Tonolli, 50, 28922 Pallanza, Italy) and P. Cardoso (E-mail: pedro.cardoso@helsinki.fi; Tel.: +358503185685; Finnish Museum of Natural History, Pohjoinen Rautatiekatu 13, 00100 Helsinki, Finland).

ABSTRACT

Five decades ago, a landmark paper in *Science* titled *The Cave Environment* heralded caves as ideal natural experimental laboratories in which to develop and address general questions in geology, ecology, biogeography, and evolutionary biology. Although the ‘caves as laboratory’ paradigm has since been advocated by subterranean biologists, there are few examples of studies that successfully translated their results into general principles. The contemporary era of big data, modelling tools, and revolutionary advances in genetics and (meta)genomics provides an opportunity to revisit unresolved questions and challenges, as well as examine promising new

91 avenues of research in subterranean biology. Accordingly, we have developed a roadmap to
92 guide future research endeavours in subterranean biology by adapting a well-established
93 methodology of ‘horizon scanning’ to identify the highest priority research questions across six
94 subject areas. Based on the expert opinion of 30 scientists from around the globe with
95 complementary expertise and of different academic ages, we assembled an initial list of 258
96 fundamental questions concentrating on macroecology and microbial ecology, adaptation,
97 evolution, and conservation. Subsequently, through online surveys, 130 subterranean biologists
98 with various backgrounds assisted us in reducing our list to 50 top-priority questions. These
99 research questions are broad in scope and ready to be addressed in the next decade. We believe
100 this exercise will stimulate research towards a deeper understanding of subterranean biology and
101 foster hypothesis-driven studies likely to resonate broadly from the traditional boundaries of this
102 field.

103

104 *Key words:* biospeleology, cave biology, expert opinion, groundwater, horizon scanning,
105 research questions, stygofauna, troglobionts.

106

107 CONTENTS

108 I. Introduction

109 II. Horizon scanning protocol

110 (1) Initial list assembly

111 (2) Voting procedure and selection of 50 top-priority questions

112 (3) Caveats on interpretation

113 III. Summary of the horizon scan

114	IV. Adaptation
115	V. Origin and evolution
	VI. Community ecology
	VII. Macroecology and biogeography
116	VIII. Conservation
117	IX. Microbiology and applied topics
118	X. Conclusions
119	XI. Acknowledgements
120	XII. References
121	XIII. Supporting information

122

123 I. INTRODUCTION

124 In the era of the Internet, social media, and open-access mega-journals, the amount of accessible
125 scientific information is overwhelming (Landhuis, 2016; Wakeling *et al.*, 2016; Fire & Guestrin,
126 2019; Jarić *et al.*, 2020). It is estimated that more than 50 million peer-reviewed scientific papers
127 exist (Jinha, 2010) and about 1.5 million new articles are published every year (Laurance *et al.*,
128 2013). To capitalize on the volume of this information and make the most of it (e.g. Ioannidis,
129 2005; Jeschke *et al.*, 2019), it is becoming increasingly important for scientists to explore
130 effective ways to capture the latest advances in their field or related fields of research. Horizon
131 scanning – i.e. the systematic examination of information to identify emerging issues and
132 opportunities in a given research area – has become a useful tool to summarize and determine
133 research priorities and agendas (Sutherland *et al.*, 2011). The most important questions in
134 ecology (Sutherland *et al.*, 2013; McGill *et al.*, 2019), island biogeography (Patiño *et al.*, 2017),

and microbiology (Antwis *et al.*, 2017), the annual identification of emerging issues in global conservation (Sutherland *et al.*, 2020), as well as the 100 articles that every ecologist should read (Courchamp & Bradshaw, 2018), are all instructive examples where horizon scanning has successfully synthesized trends or highlighted the most promising future research avenues.

Fifty years ago, in a landmark *Science* paper titled *The Cave Environment*, Poulson & White (1969) heralded caves as ‘natural laboratories’, i.e. simplified settings that can be used to understand the principles governing the dynamics of more complex environments. Characterized by stringent environmental constraints and simple communities, subterranean habitats have been regarded as ideal systems for investigating many of the unresolved questions in ecology, biogeography, and evolutionary biology (Juan *et al.*, 2010; Sánchez-Fernández *et al.*, 2018; Mammola, 2019). Scientists have also studied subterranean organisms to understand human diseases such as autism (Yoshizawa *et al.*, 2018), diabetes (Riddle *et al.*, 2018), and cancer (Gatenby, Gillies & Brown, 2011), to investigate the engineering potential of biologically inspired materials (Lepore *et al.*, 2012), and to discover new drugs and pharmaceutical products (Cheeptham *et al.*, 2013). Others have even looked at caves through the lens of astrobiology, showing that the subterranean microbiome might hold clues to life beyond Earth (Northup *et al.*, 2011; Popa *et al.*, 2011).

Although the ‘caves as laboratory’ paradigm is often advocated by subterranean biologists, examples of studies that have successfully translated their results into general principles remain few in number. Five decades after Poulson & White (1969), subterranean biology is entering a new research era dominated by big data (Zagmajster *et al.*, 2019), modelling tools (Flôres *et al.*, 2013; Mammola & Leroy, 2018), and increasingly cheaper molecular approaches (Pérez-Moreno, Iliffe & Bracken-Grissom, 2016; Lefébure *et al.*, 2017).

Concomitantly, we are facing a global crisis that is negatively impacting subterranean biodiversity (Mammola *et al.*, 2019b; Boulton, 2020). Therefore, the time is ripe to review the outstanding challenges faced by this broad-in-scope discipline, as well as promising new research avenues where subterranean-based studies may be helpful in answering general and broadly scoped questions. Because gathering multiple views on such an extensive subject is difficult, we relied on the well-established methodology of horizon scanning to identify 50 fundamental, but unresolved questions in subterranean biology. With this intellectual exercise, we aimed to develop a roadmap that will guide future research endeavours and stimulate hypothesis-driven studies likely to resonate beyond the boundaries of this discipline.

II. HORIZON SCANNING PROTOCOL

(1) Initial list assembly

We used horizon scanning methodology (Sutherland *et al.*, 2011) and adapted the approach developed by Patiño *et al.* (2017) to identify priority research questions in subterranean biology. Survey coordinators (S.M. and P.C.) identified seven subject areas within the subterranean biology discipline (Table 1), namely: (1) Adaptation, (2) Origin and evolution, (3) Community ecology, (4) Macroecology and biogeography, (5) Conservation biology, (6) Microbiology and applied topics, and (7) Other topics. We included the latter subject area to cover additional topics or ideas that departed from the six core subject areas and may have been overlooked. For each subject area, survey coordinators invited a senior researcher (highlighted with asterisks in Table 1) to act as panel coordinator, with the task of establishing an international panel of experts to identify and formulate a set of fundamental questions. Each panel coordinator selected and invited three or four members to join their panel, which included at least one early-career

181 scientist (i.e. a postdoc or researcher with less than 10 years of experience) to obtain a multi-
182 generational perspective on the different topics. Survey coordinators encouraged panel members
183 to consult broadly with colleagues and select additional researchers to join their panels if deemed
184 important in providing complementary expertise. In assembling the panels, our goal was to
185 maximize multidisciplinary, while ensuring that research interests within the seven panels
186 covered a broad array of geographic areas, model organisms, and networks of international
187 collaborators. Members of each panel identified at least 20 questions that they viewed as
188 fundamental within their subject area and thus likely to advance the field significantly.

189 In total, we assembled 258 questions, which were screened for duplication or ambiguity
190 by the survey coordinators. In this phase, survey coordinators purged most subterranean-specific
191 jargon from questions and homogenized wording to ensure that all questions were presented in a
192 clear and straightforward manner. Therefore, throughout the survey we operated under the
193 assumption that all questions were characterized by a similar degree of readability (Plavén-
194 Sigray *et al.*, 2017). After the cleaning procedure and removal of duplicate questions, we
195 assembled a final list of 211 survey questions (hereafter ‘List #1’). In assembling List #1, we
196 subsumed questions identified by the panel focusing on ‘Other topics’ into the six main subject
197 areas.

198 199 **(2) Voting procedure and selection of 50 top-priority questions**

200 We subjected List #1 to an initial round of online voting by all panel members (Survey #1) to
201 select the most voted 20 questions for each of the six subject areas (Fig. 1). Voting was a binary
202 choice, whereby participants scored each question as either of ‘major’ or ‘minor’ importance.
203 We randomized question order for each participant. We repeated this voting protocol in all

subsequent online surveys. Each panel member voted on all questions irrespective of subject area, although votes by panelists on their subject area were disregarded in the final ranking of Survey #1. As a result, survey coordinators culled List #1 to the 120 most-voted questions (20 questions from each of six subject areas), referred to as List #2, thus balancing the number of questions in subsequent voting rounds.

We then subjected List #2 to online voting (Survey #2) by inviting a broad community of subterranean biologists including *ca.* 170 members of the International Society on Subterranean Biology (ISSB), *ca.* 50 members of the European Cave Organism Network, *ca.* 100 members of the Anchialine mailing list, as well as other working groups and email listservs related to subterranean biology that we could identify (e.g. national biospeleological groups). Note that members of these different groups often overlapped and some of the emails were no longer active. We estimated that Survey #2 reached an upper boundary of between 200 and 250 unique recipients. Of these, 133 recipients completed the online survey.

At the end of Survey #2, we gave participants the opportunity to submit one additional question if they felt this question was missing from List #2. Thus, 25 additional questions were added to the third list of questions (List #3). Questions in List #3 were voted on by all panel members (Survey #3), and then ranked (by percentage of ‘major importance’ votes per question) together with the 120 questions from List #2. Finally, we selected the highest ranking questions to assemble a list of 50 top-priority questions.

(3) Caveats on interpretation

Some general caveats should be recognized when interpreting the results of any horizon scanning survey (e.g. Sutherland *et al.*, 2011, 2013; Seddon *et al.*, 2014; Patiño *et al.*, 2017). First, the

background knowledge and intellectual passions of the experts involved may introduce subjectivity in the formulation of the initial list of topics and questions. Second, subjectivity likely plays a role throughout the voting process, as any voting outcome may be affected by the interests of a particular group of participants. In our case, potential biases in the composition of subterranean biologists sampled may have influenced the final selection of the top-priority questions to an extent difficult to quantify precisely. For example, questions related to microbiology received the lowest share of ‘major importance’ votes (mean \pm SD: 0.69 ± 0.01). It is understood that microbiology topics are not less important or timely, it is simply that microbiologists are probably underrepresented in the subterranean biology community. Also, an imbalance in the expertise of participants may explain the substantial difference in how the highest priority questions were parsed across the six subject areas – from four in ‘Community ecology’ to 12 in ‘Conservation biology’.

To address these potential shortcomings, we adopted four countermeasures. First, we increased the survey audience, by addressing the questionnaire to different groups and associations of subterranean biologists. Second, we diversified the expertise of panel members by including early-stage to mid- and late-career researchers from different disciplines, research laboratories, and geographic areas. Third, we included a seventh panel (‘Other topics’) specifically to fill the gaps in the initial composition of proposed questions. Indeed, it has been argued that in horizon scanning, the initial division into subject areas may limit lateral thinking (Sutherland *et al.*, 2013). Finally, by allowing voters to suggest additional questions when voting in the survey, we were able to capture the range of priority topics better.

We are confident these practices minimized some of the biases inherent to this study. Importantly, we believe this 50 top-priority survey served to highlight some of the most timely

and challenging areas of interest in current and future research, rather than providing a comprehensive synthesis of research needs in modern subterranean biology.

III. SUMMARY OF THE HORIZON SCAN

In Survey #1, the percentage of ‘major importance’ votes ranged between 89% (top-voted question) and 4% (least-voted question). In the extended online voting (Survey #2), 133 voters participated, of which 71% identified ‘subterranean biology’ as their primary field of research. Although voters’ gender was slightly skewed toward males (76 men *versus* 57 women), deviation from the 1:1 male:female ratio was not significant ($\chi^2 = 2.71$; d.f. = 1; $P = 0.10$), indicating that our sample was not gender-biased. 45% of survey voters were experienced researchers, with an academic age of more than 10 years since they earned their PhD, while 29% were researchers within 10 years from their PhD. PhD and undergraduate students accounted for 16% of voters. The remaining 10% of participants were other professionals, such as research and field technicians or recreational cavers.

During Survey #2, participants suggested 28 additional questions; three questions were duplicates and were thus excluded. The remaining 25 questions were evaluated during Survey #3, and three made it to the 50 top-priority list. The lower threshold for questions was 67% of ‘major importance’ votes, whereas the top-voted question garnered 91% votes (Fig. 1).

In the following, we present the 50 top-priority questions in subterranean biology according to the results of Surveys #2 and #3 (the full list of questions is provided as online supporting information in Appendix S1). For clarity, questions were compiled into our six subject areas. We provide information about each question’s final rank (#) and percentage of ‘major importance’ votes received (%), and highlight the three questions suggested by the

273 Survey #2 participants with an asterisk (*). A glossary of terms is available in Table 2.

274

275 **IV. ADAPTATION**

276 Q1 – What are the drivers of adaptive evolution in caves? [#1; 91%]

277 Q2 – What are the main constraints to subterranean adaptation? [#4; 83%]

278 Q3 – What are the degrees of adaptive plasticity of organisms across different subterranean
279 environments? [#9; 78%]

280 Q4 – Which traits of subterranean organisms should be considered as ‘adaptive’? [#11; 78%]

281 Q5 – How have morphological and behavioural traits co-evolved in subterranean organisms?
282 [#14; 76%]

283 Q6 – What is the level and nature of reproductive isolation between cave and surface populations
284 and what reproductive barriers are typically involved? [#19; 75%]

285 Q7 – Do similar traits evolve repeatedly in subterranean organisms due to changes in the same
286 genes, genetic pathways, and/or developmental processes? [#23; 73%]

287 Q8 – Have subterranean species evolved a distinct set of convergent behaviours? [#26; 72%]

288 Q9 – Are there common developmental pathways that promote or constrain subterranean
289 adaptation? [#29; 72%]

290 Q10 – Do traits that constitute reproductive isolation evolve in the same way across independent
291 closely related subterranean populations or species? [#42; 70%]

292

293 The morphology of subterranean organisms, which show bizarre convergent adaptations even
294 across different animal phyla, has historically attracted the attention of generations of scientists
295 (Juan *et al.*, 2010) including Charles Darwin (1859). Therefore, it is no surprise that subterranean

biologists participating in this survey greatly valued the role of subterranean habitats as natural laboratories for the study of adaptive evolution. Ten questions focusing on adaptation were included in our top-50 list (Fig. 1).

Colonization of suitable habitat is the initial event leading to subterranean adaptation (details in Section V). Whatever the mode or pathway, colonizers often experience a significant change upon entering the subterranean environment (i.e. complete darkness), which results in visual sensory deprivation, challenges in locating mates and food, limited or modified food resources, and physical barriers to dispersal. Adaptive responses to these factors may involve the action of selection on plastic traits already existing in the colonizers (i.e. phenotypic plasticity; Bilandžija *et al.*, 2020), standing genetic variation, or new beneficial mutations. Understanding which of these environmental factors and adaptive responses play a primary role in subterranean adaptation, either acting alone or in various combinations, was the most important question (Q1) in our survey, selected by 91% of participants. Yet, given that some higher taxa are missing or understudied in caves (Culver & Pipan, 2019), it remains unclear what are the main constraints to subterranean adaptation (Q2) and whether specific exaptations facilitate successful colonization events (see also Q11 in Section V). Resolving how many phenotypes of subterranean dwellers depend on genetic and developmental constraints (Q9), or reflect entrapment at local peaks in adaptive landscapes or recent invasions with insufficient time for selection to alter traits, is one of the future challenges for evolutionary biologists.

Additional high-priority questions were focused on subsequent refinements of the initial adaptive responses, such as the repertoire of adaptive plasticity (Q3), the degree to which pre-existing genetic variation contributes to subterranean phenotypes, and which traits of subterranean organisms can be considered as adaptive (Q4). Historically, reduction or loss of

319 traits such as eyes and pigmentation was thought to be driven by random mutations and genetic
320 drift or by natural selection, either directly or indirectly. This controversy has continued to the
321 present, with strong adaptationist (Carlini & Fong, 2017) and non-adaptationist (Wilkens &
322 Strecker, 2017) viewpoints. Depending on the species or ecological context, it is possible that all
323 of these mechanisms have roles in subterranean adaptation. Resolving this debate will require
324 explanations at the molecular, cellular, and developmental levels in multiple lineages (Jeffery,
325 2005), and the integration of this information to infer whether convergent traits evolve repeatedly
326 in subterranean animals due to changes in the same or different genes, genetic pathways, and
327 developmental processes (Q7). Answers to all these questions will contribute to our
328 understanding concerning why some species adapt rapidly and evolve when facing new
329 environmental conditions, inside or outside caves, which is a critical question given global
330 climate change (Walther *et al.*, 2002). In turn, this could provide insights about adaptive
331 processes occurring in other ecological settings with a similar set of environmental conditions
332 (e.g. permanent darkness, constancy in climatic conditions, food scarcity), such as deep-sea
333 habitats (Trontelj, Borko & Delić, 2019; Mammola, 2020).

334 Once survival in a subterranean habitat is ensured, the successful colonizers are subject to
335 adaptive morphological and behavioural (co-)evolution (Q5). Many behavioural changes are
336 probably influenced by the essential requirements of finding food and mates in darkness, and
337 may be convergent across different subterranean lineages (Q8). Also, some subterranean animals
338 suddenly attain a new status at the top trophic level and predator release occurs. For example, in
339 the Mexican tetra, *Astyanax mexicanus* (De Filippi) (Actinopterygii: Characidae), the workhorse
340 of adaptive evolution studies in caves (Jeffery, 2009; Wilkens & Strecker, 2017; Torres-Paz *et*
341 *al.*, 2018), this new ecological status of an apex predator facilitated the evolution of a range of

behaviours that may not be sustainable in a predator-limited surface environment (Yoshizawa *et al.*, 2010; Hyacinthe, Attia & Rétaux, 2019).

Most subterranean organisms may also face subsequent invasions of their habitats by new colonizers, of both former surface-dwelling conspecifics (if they are still extant) and other competing species (e.g. Howarth *et al.*, 2007; Wynne *et al.*, 2014). Therefore, to understand subterranean adaptations fully, it is crucial to explore the degree and nature of reproductive isolation between the subterranean-adapted lineages and invading surface conspecifics (Q6). The majority of subterranean animals probably arose through the process of ecological speciation in which reproductive isolation evolved as a response to divergent selection between environments (Niemiller, Fitzpatrick & Miller, 2008; Mammola *et al.*, 2018). Thus, many subterranean adaptations should at least indirectly favour non-random mating between individuals of the derived subterranean and ancestral surface populations. Understanding this will help to address whether traits that constitute reproductive isolation evolve in the same way in independent closely related subterranean populations or species (Q10), and therefore whether and how often parallel speciation occurs in the subterranean realm. Ultimately, this would shed new light concerning the intriguing hypothesis on the predictability of evolution (Blount, Lenski & Losos, 2018).

V. ORIGIN AND EVOLUTION

Q11 – Which traits present in surface species (exaptations) facilitate successful subterranean colonization and adaptation? [#12; 77%]

Q12 – How do, and which, patterns of subterranean species diversification vary across taxa and habitats? [#13; 77%]

365 Q13 – What evolutionary processes most commonly triggered radiations of subterranean
366 organisms? [#15; 76%]

367 Q14 – Do subterranean organisms lack genetic variation and thus the ability to adapt to a
368 changing environment? [#16; 75%]

369 Q15 – Does the timeline of subterranean evolution differ among taxa, types of subterranean
370 habitats, different biogeographic areas, and different ecological settings? [#22; 74%]

371 Q16 – What are the impact(s) of biotic and abiotic factors on speciation? [#28; 72%]

372 Q17 – Why are some lineages successful at colonizing subterranean habitats while others are
373 not? [#35; 71%]

374 Q18 – How old are subterranean species? [#36; 71%]

375 Q19 – The role of evolutionary processes (convergence/divergence/evolutionary
376 stasis/parallelisms) in subterranean organisms: what are the most common evolutionary
377 processes? [#40; 70%]

378 Q20 – Are shallow subterranean habitats a gateway to colonize deep zones and is the evolution
379 of deep subterranean species conditioned with a colonization of shallow and later deeper zones?
380 [#41; 70%]

381 Q21 – What is the rate of evolution of different subterranean traits and does the degree of
382 subterranean adaptation correlate with duration of subterranean inhabitation? [#44; 69%]
383

384 Subterranean animals have long interested biologists as evolutionary models. Studies of these
385 species have endeavoured to improve our understanding of evolution, its repeatability at the
386 phenotypic (Friedrich, 2013; Porter & Sumner-Rooney, 2018), physiological (Jones, Cooper &
387 Seymour, 2019), and molecular level (Leys *et al.*, 2005; Bilandžija, Četković & Jeffery, 2012;

388 Niemiller *et al.*, 2013), its reversibility (Copilaş-Ciocianu *et al.*, 2018), and the role of drift in
389 morphological changes (Martínez *et al.*, 2017; Wilkens, 2020). The eleven questions identified
390 highlight how, despite advances in the application of genetic tools and techniques in the last 50
391 years, fundamental questions regarding the origin and evolution of subterranean animals remain
392 unanswered.

393 Two high-ranked questions (Q11 and Q17) focused on the traits that enable species to
394 successfully colonize and adapt to subterranean habitats. Additional questions focused on the
395 most common evolutionary processes (Q19), and the influence of biotic and abiotic factors (Q16)
396 that lead to different patterns of diversification across subterranean lineages (Q12). Important
397 subterranean radiations are known in all major taxonomic groups (Deharveng & Bedos, 2019),
398 but only a few of them have been well documented. These include Amphipoda (Zakšek *et al.*,
399 2019), Collembola (Lukić *et al.*, 2019), and Coleoptera (Leys *et al.*, 2003; Faille *et al.*, 2010;
400 Njunjić *et al.*, 2018). Which evolutionary processes best explain these radiations remains highly
401 debated (Q13) and it would be particularly interesting to compare and contrast radiations of
402 surface-dwelling plants and animals (Gillespie *et al.*, 2020) with subterranean-adapted species to
403 determine if any universal patterns exist. For many animal groups, subterranean species are
404 commonly assumed to have evolved from surface species (Barr & Holsinger, 1985; Peck &
405 Finston, 1993), but recent phylogenetic studies suggest that this assumption may not always
406 apply (Faille *et al.*, 2010; Juan *et al.*, 2010; Leijs *et al.*, 2012). Speciation and diversification may
407 also occur within the confines of a subterranean habitat, a process referred to as ‘endogenous
408 diversification’ (Trontelj, 2019). Moreover, some phylogenetic studies suggested that
409 subterranean colonization is not an evolutionary dead end and surface species may actually arise
410 from subterranean ancestors (Prendini, Francke & Vignoli, 2010; Niemiller *et al.*, 2013; Copilaş-

411 Ciocianu *et al.*, 2018). However, cases of endogenous speciation and ‘subterranean to surface’
412 reversals are potentially confounded by extinction of surface lineages (Juan *et al.*, 2010).
413 Therefore, new approaches are needed that avoid reliance on phylogenetic methods alone to
414 improve our understanding of these patterns.

415 Genetic variation enhances the ability of species to adapt and diversify. Additionally, it
416 has been shown that some subterranean species may contain high levels of neutral genetic
417 variation (Buhay & Crandall, 2005; Guzik *et al.*, 2009), but it is still unclear whether neutral
418 mutations equates to high levels of adaptive genetic variation. This underpins the question
419 whether subterranean species lack the ability to adapt to changing environments (Q14), including
420 increasing temperatures and the introduction of new pathogens (Mammola *et al.*, 2019c). Such
421 hypotheses are obviously not exclusive to the subterranean environment. However, this
422 ecosystem does provide numerous examples of how low genetic variation was hypothesized to
423 be related to low adaptive capacity, a phenomenon more common underground than at the
424 surface (Konec *et al.*, 2015; Lefébure *et al.*, 2017; Fumey *et al.*, 2018).

425 Understanding the timeline and direction of subterranean evolution, as well as the age of
426 subterranean species, featured prominently in several questions (Q15, Q18, Q20, Q21).
427 Advances in molecular clock calibration (Drummond *et al.*, 2006) and genomic analyses (Pérez-
428 Moreno *et al.*, 2016) are considerably promising and permit the development of robust time trees
429 (Pons *et al.*, 2019). However, these analyses are limited by the availability of extant and fossil
430 taxa and the extinction of surface relatives; the latter makes it difficult to pinpoint the initial
431 colonization time of a subterranean habitat by a given species. This is particularly important for
432 ancient lineages of specialized subterranean organisms with marine origin, which often lack
433 surface-dwelling relatives and/or show low levels of fossilization (Pérez-Moreno *et al.*, 2016).

434 This is unfortunate because many of these basally branching lineages are required to reconstruct
435 trait evolution of major animal lineages (e.g. Johnson *et al.*, 2012; Khodami *et al.*, 2017; Lozano-
436 Fernandez *et al.*, 2019).

437 The genetic basis underlying evolution of subterranean traits, and how they are shaped by
438 natural selection and/or neutral processes, are key factors in determining rates of subterranean
439 evolution (Q21). Considerable advances have been made through the study of model
440 subterranean species, especially *Astyanax mexicanus* and the freshwater isopod *Asellus aquaticus*
441 (L.) (Protas & Jeffery, 2012). These species have several independent and recently evolved
442 subterranean populations, as well as extant surface populations, which can be hybridized in the
443 laboratory. Their features allow for the dissection of genes and mutations responsible for traits
444 related to subterranean life and provide information on the processes (e.g. selection or neutral
445 evolution) that shape their evolution. The role of neutral processes in the evolution of
446 subterranean animals has also been explored using alternative model systems (e.g. dytiscid
447 beetles and amblyopsid cavefishes). In both cases, species have been evolving underground for
448 millions of years, which is sufficient to enable the fixation of deleterious mutations in genes
449 under relaxed selection (Niemiller *et al.*, 2013; Tierney *et al.*, 2018). These model organisms
450 offer great potential to investigate major questions on the origin and evolution of subterranean
451 animals using comparative genomics, and thus may provide insights for similar processes in
452 other, non-subterranean, settings.

VI. COMMUNITY ECOLOGY

Q22 – What are the main ecological and ecosystem services provided by subterranean populations and communities? [#20; 75%]

Q23 – What are the key food-web processes influencing subterranean community dynamics? [#24; 73%]

Q24 – How do stochastic events interact with long-term trends in subterranean ecosystems? [#30; 72%]

Q25 – How do basic life-history characteristics differ among subterranean communities and between subterranean and surface communities? [#33; 71%]

Subterranean habitats are well-suited systems to address general problems in community ecology (Mammola, 2019). Foremost, caves are often semi-closed environments extensively replicated across the Earth (Culver, 1970; Culver & Pipan, 2019; Itescu, 2019; Mammola, 2019). Second, subterranean communities generally exhibit lower diversity and abundance of organisms than surface ones and are characterized by a bottom-truncated functional diversity (Gibert & Deharveng, 2002), allowing us to disentangle the effect of abiotic conditions and biotic interactions in filtering species possessing specific traits within the community (Cardoso, 2012). Third, caves have some conspicuous environmental gradients from the surface towards the subsurface (Howarth, 1982; Tobin, Hutchins & Schwartz, 2013; Mammola *et al.*, 2019d), offering a mosaic structure of subterranean microhabitats defined by distinct habitat-filtering properties (Trontelj, Blejec & Fišer, 2012; Mammola *et al.*, 2020).

Four questions in community ecology made it to the top-50 list. This result reflects a general trend in subterranean biology, where researchers have primarily focused on caves as

476 model systems for evolutionary studies (Juan *et al.*, 2010), and secondarily used caves as
477 convenient settings to address fundamental ecological questions (Mammola, 2019). Yet, these
478 four questions fell within general and timely areas of current ecological research (see Sutherland
479 *et al.*, 2013).

480 The top-ranked question underscored the importance of services provided to humans by
481 subterranean species and ecosystems (Q22), rather than on theoretical aspects of community
482 ecology. Examples of ecosystem services provided by subterranean ecosystems include
483 pollination, seed dispersal, and agricultural pest control by bats (Kunz *et al.*, 2011; Medellin,
484 Wiederholt & Lopez-Hoffman, 2017), provision of clean water (Griebler & Avramov, 2015),
485 serving as a source for new pharmaceutical products (Cheeptham *et al.*, 2013), and even cheese
486 production (Ozturkoglu-Budak *et al.*, 2016). While services with direct benefit to humans have
487 received some attention, values provided by subterranean ecosystems extend far beyond direct
488 human needs. In light of emerging conservation issues associated with subterranean ecosystems
489 (Mammola *et al.*, 2019b), investigating ecological services and links between above- and below-
490 ground diversity in ecosystem functioning is crucial.

491 Two questions called for more research into life-history characteristics (e.g. growth rates,
492 age and size at sexual maturity, longevity, and survival rates; Q25) and food-web specificities of
493 subterranean communities (Q23). Interactions among life-history traits determine the fitness of
494 each population, while interactions between populations and the environment dictate the
495 distribution of species (Sterns, 1992). Only a few studies have described life histories of
496 subterranean species, and this is partially explained by the challenges of captive breeding and the
497 technical problems and effort necessary to conduct *in situ* comprehensive studies (Vonk &
498 Nijman, 2006; Voituron *et al.*, 2011; Venarsky, Huryn & Benstead, 2012; Riesch *et al.*, 2016;

Simon *et al.*, 2017). Consequently, the lack of knowledge on cave species traits limits our understanding of evolutionary and ecological processes occurring in subterranean ecosystems.

Energy limitation is considered a primary mechanism influencing both evolutionary and ecological processes in subterranean environments (Venarsky & Huntsman, 2018). However, a more nuanced understanding of subterranean food-web dynamics (Q23) will require other research actions, including to (i) understand the spatial and temporal dynamics of energy resources; (ii) compare resource quality with consumers' physiological requirements; and (iii) compare consumption rates with resource availability in subterranean habitats with different environmental conditions (e.g. terrestrial *versus* aquatic, fresh *versus* salt water, and detrital *versus* chemolithoautotrophic food webs).

Finally, understanding the role of stochastic events in caves was highlighted as a deficient area in community ecology (Q24). Given that these events are increasing in frequency amid the environmental crisis of the new millennium (Rahmstorf & Coumou, 2011), the study of stochastic phenomena has emerged as a central topic in ecology (Scheffer *et al.*, 2001). Recent papers used groundwater crustaceans to elucidate some of the mechanisms by which earthquakes affect the composition and structure of biological communities (Galassi *et al.*, 2014; Fattorini *et al.*, 2017; Fattorini, Di Lorenzo & Galassi, 2018; Morimura *et al.*, 2020). Additional studies have focused on the effect of other events, such as heavy precipitation (Calderón-Gutiérrez, Sánchez-Ortiz & Huato-Soberanis, 2018) and flooding (Pacioglu *et al.*, 2019). Although it may seem counterintuitive to study stochastic environmental shifts in caves, as they have been traditionally perceived as stable ecosystems, these examples show how caves may represent promising model systems for quantifying the impacts of abrupt environmental shifts driving ecosystem evolution (Mammola, 2019).

VII. MACROECOLOGY AND BIOGEOGRAPHY

Q26 – What drives subterranean patterns of phylogenetic and functional diversity? [#21; 75%]

Q27 – Would the use of novel molecular methods (e.g. metabarcoding, environmental DNA) provide new insights on subterranean biodiversity patterns and affect known patterns? [#27; 72%]

Q28 – What is the species richness pattern of subterranean organisms globally? [#31; 72%]

Q29 – What factors drive the relative importance of speciation, extinction, and dispersal in shaping subterranean diversity patterns across regions? [#34; 71%]

Q30 – Are current subterranean biodiversity patterns best explained by history of colonization of surface ancestors or by *in situ* speciation and dispersal in subterranean habitats? [#39; 70%]

Q31 – How can sampling effort be standardized so that comparisons of species richness are unbiased? [#43; 69%]

Over the last 20 years, research in subterranean ecology is shifting from local to landscape studies aiming to document and understand biodiversity patterns at regional to global scales (Zagmajster *et al.*, 2019). This transition is not without difficulties, as it requires linking biodiversity patterns to eco-evolutionary processes with little to no possibility for manipulative experiments. Six questions in ‘Macroecology and biogeography’ were identified in the top-50 list (Fig. 1). These questions mirror the main challenges faced when documenting and understanding broad-scale biodiversity patterns at the surface. The first challenge is assembling the data required to bring out the characteristic features of biodiversity patterns at such broad scales, while ensuring these patterns are not biased by sampling effort (Q28, Q31). Secondly, to combine multiple sampling techniques, species identification methods (e.g. morphological and

DNA-based identification), and biodiversity metrics (e.g. alpha, beta, and gamma diversity) in a meaningful way to elucidate the many facets of biodiversity patterns (e.g. taxonomic, phylogenetic, and/or functional diversity; Jarzyna & Jetz, 2016) (Q27, Q26). Lastly, the relative contributions of different evolutionary processes (Q29) and diversification hypotheses (Q30) in shaping biodiversity patterns should be fully examined.

The publication of global subterranean diversity maps and databases is a recent phenomenon (Culver & Pipan, 2019; Zagamajster *et al.*, 2019). While diversity maps are informative as they portray differences in species richness among regions or countries, we still lack global maps showing species richness for spatial units of equal area [but see Zagamajster, Culver & Sket (2008), Niemiller & Zigler (2013), and Eme *et al.* (2015) for examples of regional- and continental-scale diversity maps]. Several approaches have been developed to minimize differences in species richness due to sampling bias (Q31). This issue is particularly germane to difficulties in sampling subterranean habitats. For example, sampling protocols were typically standardized among sites and completeness of species inventories were assessed using accumulation and rarefaction curves (Zagamajster *et al.*, 2008; Dole-Olivier *et al.*, 2009; Wynne *et al.*, 2018). Also, observed species richness patterns were tested for robustness using species richness estimators (Zagamajster *et al.*, 2014), or complemented with species richness predictions modelled from environmental data (Mokany *et al.*, 2019).

Beyond accounting for sampling biases, molecular methods are increasingly useful in understanding subterranean biodiversity patterns (Q27). For example, a recent study comparing latitudinal patterns of crustacean species range size obtained from morphology- and DNA-based species delimitation showed that the pattern of increasing median range size at higher latitudes was more evident when delimiting species with DNA (Eme *et al.*, 2018) (Fig. 2). As sequencing

becomes increasingly applied to subterranean taxa, environmental DNA sampling and monitoring may be also used to detect these species in areas difficult to access (Gorički *et al.*, 2017; Niemiller *et al.*, 2018), thus resulting in more accurate maps of their distributions. To our knowledge, patterns of phylogenetic and functional diversity at continental to global scales have not been documented for any subterranean taxon (Q26), despite the growing knowledge of phylogenetic relationships and species traits (Morvan *et al.*, 2013; Fernandes, Batalha & Bichuette, 2016; Fišer *et al.*, 2019; Mammola *et al.*, 2020). Documenting these patterns will further underscore the relative importance of dispersal, extinction, and different speciation modes in shaping geographic variation of species richness. Given the differences in global diversity patterns between subterranean and surface habitats, comparing the two systems might help further to elucidate the key drivers of diversity.

Recent macroecological studies have shown that historical climatic variability, spatial heterogeneity, and energy contribute to species richness patterns of subterranean taxa in Europe. However, the contributions of these factors vary regionally and across taxa (Eme *et al.*, 2015; Bregović & Zgamažster, 2016; Bregović, Fišer & Zgamažster, 2019; Mammola *et al.*, 2019a). At a landscape scale, linking environmental factors with speciation, extinction, and dispersal dynamics (Q29), as well as diversification processes (Q30), remains challenging and requires the use of phylogenetic methods and a large number of specimens for DNA analysis (Stern *et al.*, 2017). Yet phylogenetic methods encompass uncertainties that are highly sensitive to sampling bias and the confounding effect of extinction, both obscuring the inference of transitions to subterranean life. To ameliorate this, genes that lose their function soon after the transition should be used (Lefébure *et al.*, 2017) (see also Section V).

VIII. CONSERVATION

Q32 – How does climate change affect subterranean-adapted organisms? [#2; 84%]

Q33 – What are the effects of pollution on subterranean-restricted microorganisms, arthropods, and vertebrates? [#3; 84%]

Q34 – What is the impact of above-ground disturbance on subterranean environments and their fauna? [#5; 82%]

Q35 – How can we evaluate the ecological status of subterranean ecosystems? [#6; 80%]

Q36 – How can we protect subterranean-adapted species from invasive species? [#7; 80%]

Q37 – How can we combine policy, education, research, and management to safeguard subterranean biodiversity effectively? [#8; 80%]

Q38* – What factors determine the size and location of effective protected areas in subterranean environments? [#10; 78%]

Q39* – How can we effectively involve governments and key stakeholders in the conservation of caves and other subterranean systems? [#17; 75%]

Q40 – What would be the best monitoring protocols to quantify long-term changes in the distribution and abundance of subterranean invertebrates? [#18; 75%]

Q41 – How do we address the lack of knowledge (biodiversity shortfalls) about the biology of subterranean species to enhance proper conservation measures? [#25; 73%]

Q42 – Can subterranean-adapted organisms be used as bioindicators of the health of subterranean ecosystems? [#45; 69%]

Q43 – How does the use of caves by humans (e.g. tourism, religious, therapeutic, and recreational activities) affect subterranean ecosystems? [#48; 68%]

Ecosystems are experiencing biodiversity loss at an unprecedented rate worldwide (Barnosky *et al.*, 2011; Dirzo *et al.*, 2014; IPBES, 2018; Cardoso *et al.*, 2020). Thus, conservation and management of cave biological diversity is of the utmost concern among subterranean biologists (Mammola *et al.*, 2019b). Conservation questions comprised most of the questions (24%) in our top-50 list (Fig. 1). Of these, 10 questions were part of the initial List #1, while two additional questions were suggested by survey participants. Three questions (Q32, Q33, and Q36) highlighted three of the greatest threats to biodiversity worldwide – climate change (Ripple *et al.*, 2019), pollution (Ripple *et al.*, 2017), and invasive alien species (Pyšek *et al.*, 2020) – whose effects are pervasive also underground (Mammola *et al.*, 2019b). Additional questions were centred on the impacts of above-ground disturbance (Q34) and human activities (Q43) on subterranean habitats. All these threats can be combined and described as ‘habitat loss and degradation’, which is one of the most important drivers of biodiversity loss globally (IPBES, 2018). Subterranean habitat loss and degradation is primarily due to surface activities, such as agricultural expansion and intensification, urbanization, and mining activities (Reboleira *et al.*, 2013; Mammola *et al.*, 2019b; Castaño-Sánchez, Hose & Reboleira, 2020). Human activities inside caves may also constitute localized threats, with recreational use and tourism activities being of particular concern (Fernandez-Cortes *et al.*, 2011; Faille, Bourdeau & Deharveng, 2015). In certain areas, people are even poaching rare invertebrate species for private collections (Simičević, 2017), as in the discussed case of *Anopthalmus hitleri* Scheibel (Coleoptera: Carabidae) (Berenbaum, 2010).

Evaluating, understanding, and mitigating these threats are primarily hampered by our scarce knowledge of subterranean organisms’ biology (Q41), especially life-history traits (see Q25 in Section VI). Understanding changes in species’ abundance and distribution will be

637 crucial to halting biodiversity loss in subterranean habitats. Studies aimed at identifying
638 bioindicator species (Q42) to help bolster long-term monitoring programs (Q40) are needed.
639 Additionally, improved sampling procedures and characterizing cave communities in previously
640 undocumented areas would both enhance our knowledge of subterranean biodiversity (Mammola
641 *et al.*, 2019b) and improve the effectiveness of conservation measures (Q41).

642 Furthermore, it is crucial to adopt innovative approaches to safeguard subterranean
643 biodiversity (Q37), as well as to determine the size and location of effective protected areas
644 (Q38). Standardized systematic sampling techniques have been applied to terrestrial (Wynne *et*
645 *al.*, 2018, 2019) and aquatic subterranean invertebrate species (Dole-Olivier *et al.*, 2009); to be
646 optimally beneficial to conservation and monitoring, these techniques will need to be further
647 scrutinized across a large breadth of taxa and systems. Recently, a cave vulnerability assessment
648 protocol has been developed for bat cave roosts (Tanalgo, Tabora & Hughes, 2018) and, if
649 refined, would hold promise for use with other subterranean animals.

650 Protected areas are the most crucial measure to safeguard specific subterranean habitats
651 and the sensitive animal populations they often support (Q38). Indices have been developed for
652 site selection and conservation prioritization (e.g. Borges *et al.*, 2012; Rabelo, Souza-Silva &
653 Ferreira, 2018; Strona *et al.*, 2019; Fattorini *et al.*, 2020) which are often based on
654 complementarity, flexibility, and irreplaceability principles (Michel *et al.*, 2009). Yet, rigorous
655 geospatial analysis is still rarely applied when the extents of protected areas are being
656 determined. Further considerations should include managing lands upslope from caves or entire
657 watersheds supporting sensitive subterranean habitats. If a species-level approach is taken for
658 establishing a protected area, it would be reasonable to protect the land at the hydrogeologic unit
659 (i.e. watershed or karst/volcanic unit) level – as animals are expected to use mesocaverns or

unconsolidated sediments for dispersal (Howarth, 1983; Malard *et al.*, 2017; Trontelj, 2019).
Importantly, such an approach should be based on the most accurate estimation of the relevant
animal's distributional range.

While effective legislation and/or management plans exist for some subterranean species
and some regions of the world, overall management policies for most regions of speleological
importance are lacking (Q39). Only a few countries have national cave protection laws. For
example, the United States Federal Cave Protection Act of 1988 has been used as a tool to
manage caves on federally owned lands, while Brazil requires geological and biological
assessments of caves and stipulates mitigation of any human activities that may negatively
impact cave natural resources. In any case, to be fully operational, such legislative and
management tools need to be based on the best available science including a comprehensive
knowledge of fauna distribution (Brooks, Da Fonseca & Rodrigues, 2004; Samways *et al.*, 2020)
and traits of the species of concern (Chichorro, Juhlén & Cardoso, 2019; Fattorini *et al.*, 2020).
Importantly, management plans will require both financial, governmental, and local community
support for their implementation. Unfortunately, most countries lack the capacity or legislation to
protect and conserve sensitive subterranean resources.

IX. MICROBIOLOGY AND APPLIED TOPICS

Q44 – What is the role of Bacteria, Archaea, fungi, and viruses in nutrient cycling in
subterranean systems? [#32; 71%]

Q45 – How adaptable are cave microorganisms to changing environmental conditions (e.g.
climate change)? [#37; 70%]

Q46 – How do other organisms (humans and other animals), and their activities (e.g. visiting

683 humans and global climate change) influence cave microbiome diversity patterns? [#38; 70%]
684 Q47 – How does the range of energy sources and quantity influence the diversity of subterranean
685 microbiota? [#46; 68%]
686 Q48 – What are the limiting nutrients for subterranean microbiota and how do they affect overall
687 subterranean microbial diversity? [#47; 68%]
688 Q49 – How do subterranean microorganisms cycle key elements – nitrogen, iron, carbon, sulfur,
689 and phosphorus? [#49; 67%]
690 Q50* – What is the role of microorganisms in cave-formation processes (speleogenesis) in
691 subterranean environments? [#50; 67%]
692
693 Without a doubt, topics such as adaptation, origin and evolution, community dynamics, and
694 biogeographic distribution patterns are similarly important and actively targeted in microbial
695 ecology (Antwis *et al.*, 2017). However, research in macroecology and microbial ecology is
696 often conducted separately rather than hand-in-hand. For nearly 200 years, subterranean
697 ecosystems have been studied from a macroscopic perspective. Subterranean microbiological
698 research is a relatively new discipline with most research having been conducted since the
699 middle of the last century (Griebler & Lueders, 2009). A modern ecosystem approach to
700 subterranean biota requires consideration across all trophic levels and scales (Hershey & Barton,
701 2019), especially since the 1980s, when the first cave ecosystems fully sustained by *in situ*
702 chemosynthetic primary production were discovered (Sarbu, Kane & Kinkle, 1996; Kumaresan
703 *et al.*, 2014).
704 The seven questions on the top-50 list address general problems that have been frequently
705 examined for various subterranean ecosystems, such as alluvial aquifers, however, less

systematically for cave environments. Three questions focused on the active role of microorganisms in nutrient cycling (Q44, Q49) and how nutrient limitations influence microbial diversity (Q48). Although we know that microbes rule the subsurface in terms of element cycles (Ortiz *et al.*, 2014; Kimble *et al.*, 2018) and constitute the basis of the food web, we still lack detailed information on conversion rates and growth kinetics. In addition, subterranean organisms often persist with limited energy resources. Thus, understanding their specific adaptations would help advance our understanding of adaptive strategies for microorganisms in other ecosystems (e.g. mountain-summit and deep-sea habitats). Additionally, the role of viruses, which only recently has been recognized as ‘tremendous’ for groundwater ecosystems (Griebler, Malard & Lefébure, 2014), has not been investigated for terrestrial subterranean systems (Q44).

Two questions further addressed the resistance and resilience of cave microbial communities to disturbance from changes in environmental conditions (Q45) (Cavicchioli *et al.*, 2019), and the impacts of other organisms (in particular, humans; Moldovan *et al.*, 2020; Martínez *et al.*, 2020) on microbial diversity (Q46). These questions also were related to conservation issues from a microbiological perspective. The adverse impacts of the fungus *Pseudogymnoascus destructans* that causes white-nose syndrome in North American bats is a prominent example. To date, *P. destructans* occurs in 38 U.S. states and seven Canadian provinces (see <http://www.whitenosesyndrome.org>), which raises serious concerns for the conservation of hibernating bat species and the ecosystem services they provide (Kunz *et al.*, 2011; Boyles *et al.*, 2011; Medellín *et al.*, 2017; Mammola *et al.*, 2019b). The fungus is an opportunistic environmental pathogen, which can remain in the subterranean environment and contribute to the cave microbiome even in the absence of its host (Lorch *et al.*, 2013).

It has been hypothesized that microbial communities with high diversity and functional

redundancy do not select for ecosystems poor in energy and stable in environmental conditions (Griebler & Lueders, 2009). Thus, the introduction of novel species may have a destabilizing effect on a cave's biological equilibrium (Q46). The same is true for the introduction of contaminants, such as organic compounds and nutrients that provide additional energy. We are only beginning to understand whether and how energy–diversity relationships known from macroecology apply to complex natural bacterial communities (Q47). In fact, there is a growing body of evidence that diversity–productivity relationships also drive microbial communities (Smith, 2007), but this question has not been examined systematically in subterranean ecosystems yet.

Finally, Q50 points to the potential contribution of microorganisms in speleogenetic processes, such as weathering and rock formation *via* inducing precipitation. Specifically, in terms of (inorganic) carbon cycling in face of climate change, the role of microbes in the formation of caves may be of great relevance, and has yet to be fully examined.

X. CONCLUSIONS

(1) The 50th anniversary of Poulson & White's (1969) article was the perfect time to reflect on milestone scientific achievements obtained in the natural laboratories offered by caves, while also delineating the most important research priorities for years to come. We have shown how subterranean biology has contributed strongly to general scientific questions *via* the study of evolutionary and ecological processes along the vertical dimension (i.e. the evolutionary transition from the surface to the subsurface). These accomplishments resonate with the sentiments of Poulson & White (1969) and we anticipate that biologists will continue to unravel the mysteries of subterranean ecosystems and contribute to scientific knowledge more broadly,

insofar as revolutionary advances in approaches and technologies continue to foster and nurture novel paradigms.

(2) There is a significant lack of knowledge concerning eco-evolutionary processes underlying biodiversity patterns along the horizontal gradient (i.e. within subterranean habitats). This is largely driven by a paucity of functional ecology studies, the weakness of trait-based approaches (Cardoso, 2012; Fernandes *et al.*, 2016; Fišer *et al.*, 2019; Mammola *et al.*, 2020), and the lack of robust systematic sampling techniques for most taxonomic groups (Wynne *et al.*, 2019).

Bridging these gaps will significantly influence how we address and prioritize future research on the conservation and ecosystem services of subterranean habitats (e.g. Fattorini *et al.*, 2020), as emphasized by the large number of unresolved questions in conservation biology (representing nearly 25% of the top-50 list).

(3) We also invite scientists to redouble their efforts to understand the diversity of subterranean life across all its components, with a special focus on linking macroscopic and microbial ecology (Foulquier *et al.*, 2011; Mermillod-Blondin, 2011). This will enable us to achieve a mechanistic understanding of subterranean eco-evolutionary processes and ecosystem function. This information will be critical in guiding future policy decisions as human activities and global environmental change increasingly impact and strain the subterranean realm.

(4) There is a concern that simple voting exercises such as this one may favour general over specific questions. Perhaps as a result of this bias, some of the top-voted questions appear to be broad in scope (e.g. Q1, Q2, and Q32). While these questions were able to capture important general lines of inquiry, specific questions may be more useful for setting applied agendas.

Therefore, we invite interested readers to consult Appendix S1, which contains our complete list of 120 questions.

(5) While the ‘caves as laboratory’ paradigm is an effective way to frame broadly scoped studies, we recognize the top-50 list of questions primarily pertains to unresolved issues within the borders of subterranean biology. Yet subterranean habitats offer much more. Deep subterranean habitats are one of the few natural systems defined by highly stable and homogenous climatic conditions tantamount to those maintained in a laboratory (Sánchez-Fernández *et al.*, 2018). These systems have an island-like nature (Itescu, 2019), and often support communities characterized by highly specialized organisms interacting in simplified ecological networks (Mammola, 2019). By extension, a robust understanding of these rather simplified settings may enable researchers to disentangle the complexities of more diverse systems (e.g. deep-sea habitats).

(6) Ultimately, all these features point at subterranean ecosystems as ideal settings in which to tackle general questions. We strived to provide examples of how some of our survey questions may aid in addressing non-cave specific agendas. Our hope is that this horizon scan exercise both underscores the importance of caves for addressing a range of eco-evolutionary questions, as well as stimulates researchers to redouble their efforts to address some of these lingering questions in subterranean biology.

XI. ACKNOWLEDGEMENTS

A special thanks to our colleagues around the world who participated in the online survey and made this paper possible. Thanks to Alison Cooper, Simone Fattorini and two anonymous referees for useful suggestions through the review process. S.M. was supported by the CAWEB project “*Testing macroecological theory using simplified systems*”, funded by the European Commission through Horizon 2020 Marie Skłodowska-Curie Actions (MSCA) individual

798 fellowships (Grant no. 882221). I.R.A. was supported by Portuguese funds through FCT –
 799 Fundação para a Ciência e a Tecnologia, I.P., under the Norma Transitória –
 800 DL57/2016/CP1375/CT0003. D.E. was supported by IFREMER and by the CERES “*Climate*
 801 *change and European Aquatic Resources*” project funded by European Commission through
 802 Horizon 2020 research and innovation programme (Grant no. 678193). C.F., Z.F., and M.Z. were
 803 supported by the Slovenian Research Agency (program P1-0184, project N1-0069). J.E.K. was
 804 funded by NSF awards DEB1754231 and IOS1933428, and EDGE award 1923372. F.M. was
 805 supported by the French National Research Agency projects CONVERGENOMICS (ANR-15-
 806 CE32-0005) and EUR H2O’Lyon (ANR-17-EURE-0018). A.M. was supported by the ANCAVE
 807 project “*Anchialine caves to understand evolutionary processes*”, funded by the European
 808 Commission through Horizon 2020 Marie Skłodowska-Curie Actions (MSCA) individual
 809 fellowships (Grant no. 745530). P.A.V.B. was supported by the project AZORESBIOPORTAL –
 810 PORBIOTA (ACORES-01-0145-FEDER-000072), financed by FEDER in 85% and by Azorean
 811 Public funds by 15% through Operational Program Azores 2020. A.S.P.S.R. was supported by
 812 VILLUM FONDEN (Grant no. 15471) and by a Carlsbergfondet grant (CF19-0609). T.G.P. was
 813 supported by Vale S.A. and Fundação de Amparo à Pesquisa de Minas Gerais (FAPEMIG) for
 814 individual fellowship (RDP 00092-18). T.P. was supported by the Karst Research Programme
 815 P6-0119, LifeWatch ERIC, RI-SI LifeWatch, and EU H2020 project eLTER.
 816 **Author contributions:** S.M. and P.C. conceived the idea, coordinated the survey, and curated
 817 the lists of questions. T.P., S.J.B.C., R.L.F., F.M., P.A.V.B., T.M.L., and D.C.C. coordinated
 818 research panels to identify research questions. All authors except S.M. and P.C. assembled the
 819 initial list of questions (see Table 1 for details). All authors were involved in online voting, and
 820 contributed to manuscript writing.

821

822 **XII. REFERENCES**

- 823 ANTWIS, R.E., GRIFFITHS, S.M., HARRISON, X.A., ARANEGA-BOU, P., ARCE, A., BETTRIDGE, A.S.,
824 BRAILSFORD, F.L., DE MENEZES, A., DEVAYNES, A., FORBES, K.M., FRY, E.L., GOODHEAD,
825 I., HASKELL, E., HEYS, C., JAMES, C., *ET AL.* (2017). Fifty important research questions in
826 microbial ecology. *FEMS Microbiology Ecology* **93**, 10.1093/femsec/fix044.
- 827 AUDRA, P. & PALMER, A.N. (2011). The pattern of caves: controls of the epigenic speleogenesis.
828 *Géomorphologie* **17**, 359–378.
- 829 BARNOSKY, A.D., MATZKE, N., TOMIYA, S., WOGAN, G.O.U., SWARTZ, B., QUENTAL, T.B.,
830 MARSHALL, C., MCGUIRE, J.L., LINDSEY, E.L., MAGUIRE, K.C., MERSEY, B. & FERRER,
831 E.A. (2011). Has the Earth’s sixth mass extinction already arrived? *Nature* **471**, 51–57.
- 832 BARR, T.C. & HOLSINGER, J.R. (1985). Speciation in cave faunas. *Annual Review of Ecology and*
833 *Systematics* **16**, 313–337.
- 834 BERENBAUM, M. (2010). ICE Breakers. *American Entomologist* **56**, 132–185.
- 835 BILANDŽIJA, H., ČETKOVIĆ, H. & JEFFERY, W.R. (2012). Evolution of albinism in cave
836 planthoppers by a convergent defect in the first step of melanin biosynthesis. *Evolution &*
837 *Development* **14**, 196–203.
- 838 BILANDŽIJA, H., HOLLIFIELD, B., STECK, M., MENG, G., NG, M., KOCH, A.D., GRAČAN, R.,
839 ČETKOVIĆ, H., PORTER, M.L., RENNER, K.J. & JEFFERY, W. (2020). Phenotypic plasticity as
840 a mechanism of cave colonization and adaptation. *eLife* **9**, e51830.
- 841 BLOUNT, Z.D., LENSKE, R.E. & LOSOS, J.B. (2018). Contingency and determinism in evolution:
842 replaying life’s tape. *Science* **362**, eaam5979.
- 843 BORGES, P.A.V., CARDOSO, P., AMORIM, I.R., PEREIRA, F., CONSTÂNCIA, J.P., NUNES, J.C.,

844 BARCELOS, P., COSTA, P., GABRIEL, R. & DAPKEVICIUS, M. DE L. (2012). Volcanic caves:
845 priorities for conserving the Azorean endemic troglobiont species. *International Journal of*
846 *Speleology* **41**, 101–112.

847 BOULTON, A.J. (2020). Conservation of groundwaters and their dependent ecosystems:
848 integrating molecular taxonomy, systematic reserve planning and cultural values. *Aquatic*
849 *Conservation: Marine and Freshwater Ecosystems* **30**, 1–7.

850 BOYLES, J.G., CRYAN, P.M., MCCracken, G.F. & KUNZ, T.H. (2011). Economic importance of
851 bats in agriculture. *Science* **332**, 41–42.

852 BREGOVIĆ, P., FIŠER, C. & ZAGMAJSTER, M. (2019). Contribution of rare and common species to
853 subterranean species richness patterns. *Ecology and Evolution* **9**, 11606–11618.

854 BREGOVIĆ, P. & ZAGMAJSTER, M. (2016). Understanding hotspots within a global hotspot –
855 identifying the drivers of regional species richness patterns in terrestrial subterranean
856 habitats. *Insect Conservation and Diversity* **9**, 268–281.

857 BROOKS, T., DA FONSECA, G.A.B. & RODRIGUES, A. (2004). Species, data, and conservation
858 planning. *Conservation Biology* **18**, 1682–1688.

859 BUHAY, J.E. & CRANDALL, K.A. (2005). Subterranean phylogeography of freshwater crayfishes
860 shows extensive gene flow and surprisingly large population sizes. *Molecular Ecology* **14**,
861 4259–4273.

862 CALDERÓN-GUTIÉRREZ, F., SÁNCHEZ-ORTIZ, C.A. & HUATO-SOBERANIS, L. (2018). Ecological
863 patterns in anchialine caves. *PLoS ONE* **13**, e0202909.

864 CARDOSO, P. (2012). Diversity and community assembly patterns of epigean vs. troglobiont
865 spiders in the Iberian Peninsula. *International Journal of Speleology* **41**, 83–94.

866 CARDOSO, P., BARTON, P.S., BIRKHOFFER, K., CHICHORRO, F., DEACON, C., FARTMANN, T.,

867 FUKUSHIMA, C.S., GAIGHER, R., HABEL, J.C., HALLMANN, C.A., HILL, M.J., HOCHKIRCH, A.,
868 KWAK, M.L., MAMMOLA, S., ARI NORIEGA, J., *ET AL.* (2020). Scientists' warning to
869 humanity on insect extinctions. *Biological Conservation* **242**, 108426.

870 CARLINI, D.B. & FONG, D.W. (2017). The transcriptomes of cave and surface populations of
871 *Gammarus minus* (Crustacea: Amphipoda) provide evidence for positive selection on cave
872 downregulated transcripts. *PLoS ONE* **12**, e0186173.

873 CASTAÑO-SÁNCHEZ, A., HOSE, G.C. & REBOLEIRA, A.S.P.S. (2020). Ecotoxicological effects of
874 anthropogenic stressors in subterranean organisms: a review. *Chemosphere* **244**, 125422.

875 CAVICCHIOLI, R., RIPPLE, W.J., TIMMIS, K.N., AZAM, F., BAKKEN, L.R., BAYLIS, M.,
876 BEHRENFELD, M.J., BOETIUS, A., BOYD, P.W., CLASSEN, A.T., CROWTHER, T.W.,
877 DANOVARO, R., FOREMAN, C.M., HUISMAN, J., HUTCHINS, D.A., *ET AL.* (2019). Scientists'
878 warning to humanity: microorganisms and climate change. *Nature Reviews Microbiology*
879 **17**, 569–586.

880 CHEEPHTAM, N., SADOWAY, T., RULE, D., WATSON, K., MOOTE, P., SOLIMAN, L.C., AZAD, N.,
881 DONKOR, K.K. & HORNE, D. (2013). Cure from the cave: volcanic cave actinomycetes and
882 their potential in drug discovery. *International Journal of Speleology* **42**, 35–47.

883 CHICHORRO, F., JUSLÉN, A. & CARDOSO, P. (2019). A review of the relation between species
884 traits and extinction risk. *Biological Conservation* **237**, 220–229.

885 COPILAȘ-CIOCIANU, D., FIȘER, C., BORZA, P. & PETRUSEK, A. (2018). Is subterranean lifestyle
886 reversible? Independent and recent large-scale dispersal into surface waters by two species
887 of the groundwater amphipod genus *Niphargus*. *Molecular Phylogenetics and Evolution*
888 **119**, 37–43.

889 COURCHAMP, F. & BRADSHAW, C.J.A. (2018). 100 articles every ecologist should read. *Nature*

890 *Ecology & Evolution* **2**, 395–401.

891 CULVER, D.C. (1970). Analysis of simple cave communities I. Caves as islands. *Evolution* **24**,

892 463–474.

893 CULVER, D.C. & PIPAN, T. (2019). *The biology of caves and other subterranean habitats*. Oxford

894 University Press, USA.

895 CURL, L.R. (1964). On the definition of a cave. *National Speleological Society Bulletin* **26**, 1–6.

896 DARWIN, C. (1859). *On the origin of species by means of natural selection, or the preservation of*

897 *favoured races in the struggle of life*. John Murray, London.

898 DEHARVENG, L. & BEDOS, A. (2019). Diversity of terrestrial invertebrates in subterranean

899 habitats. In *Cave Ecology* (eds T. MOLDOVAN, L. KOVÁČ & S. HALSE), pp. 107–172.

900 Springer, Cham, Switzerland.

901 DIRZO, R., YOUNG, H.S., GALETTI, M., CEBALLOS, G., ISAAC, N.J.B. & COLLEN, B. (2014).

902 Defaunation in the Anthropocene. *Science* **345**, 401–406.

903 DOLE-OLIVIER, M.J., CASTELLARINI, F., COINEAU, N., GALASSI, D.M.P., MARTIN, P., MORI, N.,

904 VALDECASAS, A. & GIBERT, J. (2009). Towards an optimal sampling strategy to assess

905 groundwater biodiversity: comparison across six European regions. *Freshwater Biology* **54**,

906 777–796.

907 DRUMMOND, A.J., HO, S.Y.W., PHILLIPS, M.J. & RAMBAUT, A. (2006). Relaxed phylogenetics

908 and dating with confidence. *PLoS Biology* **4**, e88.

909 EME, D., ZAGMAJSTER, M., DELIĆ, T., FIŠER, C., FLOT, J.-F., KONECNY-DUPRÉ, L., PÁLSSON, S.,

910 STOCH, F., ZAKŠEK, V., DOUADY, C.J. & MALARD, F. (2018). Do cryptic species matter in

911 macroecology? Sequencing European groundwater crustaceans yields smaller ranges but

912 does not challenge biodiversity determinants. *Ecography* **41**, 424–436.

913 EME, D., ZAGMAJSTER, M., FIŠER, C., GALASSI, D., MARMONIER, P., STOCH, F., CORNU, J.F.,
 914 OBERDORFF, T. & MALARD, F. (2015). Multi-causality and spatial non-stationarity in the
 915 determinants of groundwater crustacean diversity in Europe. *Ecography* **38**, 531–540.

916 FAILLE, A., BOURDEAU, C. & DEHARVENG, L. (2015). Weak impact of tourism activities on
 917 biodiversity in a subterranean hotspot of endemism and its implications for the conservation
 918 of cave fauna. *Insect Conservation and Diversity* **8**, 205–215.

919 FAILLE, A., RIBERA, I., DEHARVENG, L., BOURDEAU, C., GARNERY, L., QUÉINNEC, E. & DEUVE,
 920 T. (2010). A molecular phylogeny shows the single origin of the Pyrenean subterranean
 921 *Trechini* ground beetles (Coleoptera: Carabidae). *Molecular Phylogenetics and Evolution*
 922 **54**, 97–106.

923 FATTORINI, S., LOMBARDO, P., FIASCA, B., DI CIOCCIO, A., DI LORENZO, T. & GALASSI, D.M.P.
 924 (2017). Earthquake-related changes in species spatial niche overlaps in spring communities.
 925 *Scientific Reports* **7**, 443.

926 FATTORINI, S., DI LORENZO, T. & GALASSI, D.M.P. (2018). Earthquake impacts on
 927 microcrustacean communities inhabiting groundwater-fed springs alter species-abundance
 928 distribution patterns. *Scientific Reports* **8**, 1501.

929 FATTORINI, S., FIASCA, B., DI LORENZO, T., DI CICCIO, M. & GALASSI, D.M.P. (2020). A new
 930 protocol for assessing the conservation priority of groundwater-dependent ecosystems.
 931 *Aquatic Conservation: Marine and Freshwater Ecosystems*, doi:10.1002/aqc.3411

932 FERNANDES, C.S., BATALHA, M.A. & BICHUETTE, M.E. (2016). Does the cave environment
 933 reduce functional diversity? *PLoS ONE* **11**, e0151958.

934 FERNANDEZ-CORTES, A., CUEZVA, S., SANCHEZ-MORAL, S., CAÑAVERAS, J.C., PORCA, E.,
 935 JURADO, V., MARTIN-SANCHEZ, P.M. & SAIZ-JIMENEZ, C. (2011). Detection of human-

936 induced environmental disturbances in a show cave. *Environmental Science and Pollution*
 937 *Research* **18**, 1037–1045.

938 FIRE, M. & GUESTRIN, C. (2019). Over-optimization of academic publishing metrics: observing
 939 Goodhart's Law in action. *GigaScience* **8**, giz053.

940 FIŠER, C., DELIĆ, T., LUŠTRIK, R., ZAGMAJSTER, M. & ALTERMATT, F. (2019). Niches within a
 941 niche: ecological differentiation of subterranean amphipods across Europe's interstitial
 942 waters. *Ecography* **42**, 1212–1223.

943 FLÔRES, D.E.F.L., TOMOTANI, B.M., TACHINARDI, P., ODA, G.A. & VALENTINUZZI, V.S. (2013).
 944 Modeling natural photic entrainment in a subterranean rodent (*Ctenomys* aff. *knighti*), the
 945 Tuco-Tuco. *PLoS ONE* **8**, e68243.

946 FOULQUIER, A., MALARD, F., MERMILOD-BLONDIN, F., MONTUELLE, B., DOLÉDEC, S., VOLAT,
 947 B. & GIBERT, J. (2011). Surface water linkages regulate trophic interactions in a
 948 groundwater food web. *Ecosystems* **14**, 1339–1353.

949 FRIEDRICH, M. (2013). Biological clocks and visual systems in cave-adapted animals at the dawn
 950 of Speleogenomics. *Integrative and Comparative Biology* **53**, 50–67.

951 FUMEY, J., HINAUX, H., NOIROT, C., THERMES, C., RÉTAUX, S. & CASANE, D. (2018). Evidence
 952 for late Pleistocene origin of *Astyanax mexicanus* cavefish. *BMC Evolutionary Biology* **18**,
 953 43.

954 GALASSI, D.M.P., LOMBARDO, P., FIASCA, B., DI CIOCCIO, A., DI LORENZO, T., PETITTA, M. &
 955 DI CARLO, P. (2014). Earthquakes trigger the loss of groundwater biodiversity. *Scientific*
 956 *Reports* **4**, 6273.

957 GATENBY, R.A., GILLIES, R.J. & BROWN, J.S. (2011). Of cancer and cave fish. *Nature Reviews*
 958 *Cancer* **11**, 237.

959 GIBERT, J. & DEHARVENG, L. (2002). Subterranean ecosystems: a truncated functional
 960 biodiversity. *BioScience* **52**, 473–481.

961 GILLESPIE, R.G., BENNETT, G.M., DE MEESTER, L., FEDER, J.L., FLEISCHER, R.C., HARMON, L.J.,
 962 HENDRY, A.P., KNOPE, M.L., MALLET, J., MARTIN, C., PARENT, C.E., PATTON, A.H.,
 963 PFENNIG, K.S., RUBINOFF, D., SCHLUTER, D., *ET AL.* (2020). Comparing adaptive radiations
 964 across space, time, and taxa. *Journal of Heredity* **111**, 1–20.

965 GORIČKI, Š., STANKOVIC, D., SNOJ, A., KUNTNER, M., JEFFERY, W.R., TRONTELJ, P., PAVIC, M.,
 966 GRIZELJ, Z., NAPARUS-ALJANCIC, M. & ALJANCIC, G. (2017). Environmental DNA in
 967 subterranean biology: Range extension and taxonomic implications for *Proteus*. *Scientific*
 968 *Reports* **7**, 91–93.

969 GOULD, S.J. & VRBA, E.S. (1982). Exaptation – a missing term in the science of form.
 970 *Paleobiology* **8**, 4–15.

971 GRIEBLER, C. & AVRAMOV, M. (2015). Groundwater ecosystem services: a review. *Freshwater*
 972 *Science* **34**, 355–367.

973 GRIEBLER, C. & LUEDERS, T. (2009). Microbial biodiversity in groundwater ecosystems.
 974 *Freshwater Biology* **54**, 649–677.

975 GRIEBLER, C., MALARD, F. & LEFÉBURE, T. (2014). Current developments in groundwater
 976 ecology-from biodiversity to ecosystem function and services. *Current Opinion in*
 977 *Biotechnology* **27**, 159–167.

978 GUZIK, M.T., COOPER, S.J.B., HUMPHREYS, W.F. & AUSTIN, A.D. (2009). Fine-scale
 979 comparative phylogeography of a sympatric sister species triplet of subterranean diving
 980 beetles from a single calcrete aquifer in Western Australia. *Molecular Ecology* **18**, 3683–
 981 3698.

982 HERSHEY, O.S. & BARTON, H.A. (2019). The microbial diversity of caves. In *Cave Ecology* (eds
 983 T. MOLDOVAN, L. KOVÁČ & S. HALSE), pp. 69–90. Springer, Cham, Switzerland.
 984 HOWARTH, F.G. (1982). Bioclimatic and geologic factors governing the evolution and
 985 distribution of Hawaiian cave insects. *Entomologia Generalis* **8**, 17–26.
 986 HOWARTH, F.G. (1983). Ecology of cave arthropods. *Annual Review of Entomology* **28**, 365–389.
 987 HOWARTH, F.G., JAMES, S.A., MCDOWELL, W., PRESTON, D.J. & IMADA, C.T. (2007).
 988 Identification of roots in lava tube caves using molecular techniques: implications for
 989 conservation of cave arthropod faunas. *Journal of Insect Conservation* **11**, 251–261.
 990 HYACINTHE, C., ATTIA, J. & RÉTAUX, S. (2019). Evolution of acoustic communication in blind
 991 cavefish. *Nature Communications* **10**, 4231.
 992 IOANNIDIS, J.P.A. (2005). Why most published research findings are false. *PLOS Medicine* **2**,
 993 e124.
 994 IPBES (2018). Summary for policymakers of the thematic assessment report on land degradation
 995 and restoration of the Intergovernmental Science-Policy Platform on Biodiversity and
 996 Ecosystem Services. *Preliminary guide regarding diverse conceptualization of multiple*
 997 *values of nature and its benefits, including biodiversity and ecosystem functions and*
 998 *services (deliverable 3 (d)).*
 999 ITESCU, Y. (2019). Are island-like systems biologically similar to islands? A review of the
 1000 evidence. *Ecography* **42**, 1298–1314.
 1001 JARIĆ, I., CORREIA, R.A., BROOK, B.W., BUETTEL, J.C., COURCHAMP, F., DI MININ, E., FIRTH,
 1002 J.A., GASTON, K.J., JEPSON, P., KALINKAT, G., LADLE, R., SORIANO-REDONDO, A., SOUZA,
 1003 A.T. & ROLL, U. (2020). iEcology: harnessing large online resources to generate ecological
 1004 insights. *Trends in Ecology & Evolution* **35**, 630–639.

- JARZYNA, M.A. & JETZ, W. (2016). Detecting the multiple facets of biodiversity. *Trends in Ecology & Evolution* **31**, 527–538.
- JEFFERY, W.R. (2005). Adaptive evolution of eye degeneration in the Mexican blind cavefish. *Journal of Heredity* **96**, 185–196.
- JEFFERY, W.R. (2009). Regressive evolution in *Astyanax* cavefish. *Annual Review of Genetics* **43**, 25–47.
- JESCHKE, J.M., LOKATIS, S., BARTRAM, I. & TOCKNER, K. (2019). Knowledge in the dark: scientific challenges and ways forward. *FACETS* **4**, 423–441.
- JINHA, A.E. (2010). Article 50 million: an estimate of the number of scholarly articles in existence. *Learned Publishing* **23**, 258–263.
- JOHNSON, G.D., IDA, H., SAKAUE, J., SADO, T., ASAHIDA, T. & MIYA, M. (2012). A ‘living fossil’ eel (Anguilliformes: Protanguillidae, fam. nov.) from an undersea cave in Palau. *Proceedings of the Royal Society B: Biological Sciences* **279**, 934–943.
- JONES, K.K., COOPER, S.J.B. & SEYMOUR, R.S. (2019). Cutaneous respiration by diving beetles from underground aquifers of Western Australia (Coleoptera: Dytiscidae). *Journal of Experimental Biology* **222**, jeb196659.
- JUAN, C., GUZIK, M.T., JAUME, D. & COOPER, S.J.B. (2010). Evolution in caves: Darwin’s ‘wrecks of ancient life’ in the molecular era. *Molecular Ecology* **19**, 3865–3880.
- KHODAMI, S., MCARTHUR, J.V., BLANCO-BERCIAL, L. & MARTINEZ ARBIZU, P. (2017). Molecular phylogeny and revision of copepod Orders (Crustacea: Copepoda). *Scientific Reports* **7**, 9164.
- KIMBLE, J.C., WINTER, A.S., SPILDE, M.N., SINSABAUGH, R.L. & NORTHUP, D.E. (2018). A potential central role of Thaumarchaeota in N-Cycling in a semi-arid environment, Fort

1028 Stanton Cave, Snowy River passage, New Mexico, USA. *FEMS microbiology ecology* **94**.

1029 KONEC, M., PREVORČNIK, S., SARBU, S.M., VEROVNIK, R. & TRONTELJ, P. (2015). Parallels
 1030 between two geographically and ecologically disparate cave invasions by the same species,
 1031 *Asellus aquaticus* (Isopoda, Crustacea). *Journal of Evolutionary Biology* **28**, 864–875.

1032 KUMARESAN, D., WISCHER, D., STEPHENSON, J., HILLEBRAND-VOICULESCU, A. & MURRELL, J.C.
 1033 (2014). Microbiology of mobile cave-A chemolithoautotrophic ecosystem.
 1034 *Geomicrobiology Journal* **31**, 186–193.

1035 KUNZ, T.H., BRAUN DE TORREZ, E., BAUER, D., LOBOVA, T. & FLEMING, T.H. (2011). Ecosystem
 1036 services provided by bats. *Annals of the New York Academy of Sciences* **1223**, 1–38.

1037 LANDHUIS, E. (2016). Scientific literature: information overload. *Nature* **535**, 457–458.

1038 LAURANCE, W.F., USECHE, D.C., LAURANCE, S.G. & BRADSHAW, C.J.A. (2013). Predicting
 1039 publication success for biologists. *BioScience* **63**, 817–823.

1040 LEFÉBURE, T., MORVAN, C., MALARD, F., FRANÇOIS, C., KONECNY-DUPRÉ, L., GUÉGUEN, L.,
 1041 WEISS-GAYET, M., SEGUIN-ORLANDO, A., ERMINI, L., SARKISSIAN, C. DER, PIERRE
 1042 CHARRIER, N., EME, D., MERMILOD-BLONDIN, F., DURET, L., VIEIRA, C., *ET AL.* (2017).
 1043 Less effective selection leads to larger genomes. *Genome Research* **27**, 1016–1028.

1044 LEIJES, R., VAN NES, E.H., WATTS, C.H., COOPER, S.J.B., HUMPHREYS, W.F. & HOGENDOORN, K.
 1045 (2012). Evolution of blind beetles in isolated aquifers: a test of alternative modes of
 1046 speciation. *PLoS ONE* **7**, e34260.

1047 LEPORE, E., MARCHIORO, A., ISAIA, M., BUEHLER, M.J. & PUGNO, N.M. (2012). Evidence of the
 1048 most stretchable egg sac silk stalk, of the European Spider of the Year *Meta menardi*. *PLoS*
 1049 *ONE* **7**, e30500.

1050 LEYS, R., COOPER, S.J.B., STRECKER, U. & WILKENS, H. (2005). Regressive evolution of an eye

1051 pigment gene in independently evolved eyeless subterranean diving beetles. *Biology Letters*
1052 **1**, 496–499.

1053 LEYS, R., WATTS, C.H.S., COOPER, S.J.B. & HUMPHREYS, W.F. (2003). Evolution of
1054 subterranean diving beetles (Coleoptera: Dytiscidae: Hydroporini, Bidessini) in the arid
1055 zone of Australia. *Evolution* **57**, 2819–2834.

1056 LORCH, J.M., MULLER, L.K., RUSSELL, R.E., O’CONNOR, M., LINDNER, D.L. & BLEHERT,
1057 D.S. (2013). Distribution and environmental persistence of the causative agent of White-
1058 Nose Syndrome, *Geomyces destructans*, in bat hibernacula of the Eastern United. *Applied*
1059 *and Environmental Microbiology* **79**, 1293–1301.

1060 LOZANO-FERNANDEZ, J., GIACOMELLI, M., FLEMING, J.F., CHEN, A., VINTHER, J., THOMSEN,
1061 P.F., GLENNER, H., PALERO, F., LEGG, D.A., ILIFFE, T.M., PISANI, D. & OLESEN, J. (2019).
1062 Pancrustacean evolution illuminated by taxon-rich genomic-scale data sets with an
1063 expanded Remipede sampling. *Genome Biology and Evolution* **11**, 2055–2070.

1064 LUKIĆ, M., DELIĆ, T., PAVLEK, M., DEHARVENG, L. & ZAGMAJSTER, M. (2019). Distribution
1065 pattern and radiation of the European subterranean genus *Verhoeffiella* (Collembola,
1066 Entomobryidae). *Zoologica Scripta* **49**, 86–100.

1067 MALARD, F., CAPDERREY, C., CHURCHEWARD, B., EME, D., KAUFMANN, B., KONECNY-DUPRÉ,
1068 L., LÉNA, J.-P., LIÉBAULT, F. & DOUADY, C.J. (2017). Geomorphic influence on
1069 intraspecific genetic differentiation and diversity along hyporheic corridors. *Freshwater*
1070 *Biology* **62**, 1955–1970.

1071 MAMMOLA, S. (2019). Finding answers in the dark: caves as models in ecology fifty years after
1072 Poulson and White. *Ecography* **42**, 1331–1351.

1073 MAMMOLA, S. (2020). On deepest caves, extreme habitats, and ecological superlatives. *Trends in*

1074 *Ecology & Evolution* **35**, 469–472.

1075 MAMMOLA, S., ARNEDO, M.A., FIŠER, C., CARDOSO, P., DEJANAZ, A.J. & ISAIA, M. (2020).

1076 Environmental filtering and convergent evolution determine the ecological specialization of

1077 subterranean spiders. *Functional Ecology* **34**, 1064–1077.

1078 MAMMOLA, S., ARNEDO, M.A., PANTINI, P., PIANO, E., CHIAPPETTA, N. & ISAIA, M. (2018).

1079 Ecological speciation in darkness? Spatial niche partitioning in sibling subterranean spiders

1080 (Araneae : Linyphiidae : *Troglohyphantes*). *Invertebrate Systematics* **32**, 1069–1082.

1081 MAMMOLA, S., CARDOSO, P., ANGYAL, D., BALÁZS, G., BLICK, T., BRUSTEL, H., CARTER, J.,

1082 ĆURČIĆ, S., DANFLOUS, S., DÁNYI, L., DÉJEAN, S., DELTSHEV, C., ELVERICI, M.,

1083 FERNÁNDEZ, J., GASPARO, F., *ET AL.* (2019a). Local- versus broad-scale environmental

1084 drivers of continental β -diversity patterns in subterranean spider communities across

1085 Europe. *Proceedings of the Royal Society B: Biological Sciences* **286**, 20191579.

1086 MAMMOLA, S., CARDOSO, P., CULVER, D.C., DEHARVENG, L., FERREIRA, R.L., FIŠER, C.,

1087 GALASSI, D.M.P., GRIEBLER, C., HALSE, S., HUMPHREYS, W.F., ISAIA, M., MALARD, F.,

1088 MARTINEZ, A., MOLDOVAN, O.T., NIEMILLER, M.L., *ET AL.* (2019b). Scientists’ warning on

1089 the conservation of subterranean ecosystems. *BioScience* **69**, 641–650.

1090 MAMMOLA, S. & LEROY, B. (2018). Applying species distribution models to caves and other

1091 subterranean habitats. *Ecography* **41**, 1194–1208.

1092 MAMMOLA, S., PIANO, E., CARDOSO, P., VERNON, P., DOMÍNGUEZ-VILLAR, D., CULVER, D.C.,

1093 PIPAN, T. & ISAIA, M. (2019c). Climate change going deep: the effects of global climatic

1094 alterations on cave ecosystems. *The Anthropocene Review* **6**, 2053019619851594.

1095 MAMMOLA, S., PIANO, E., MALARD, F., VERNON, P. & ISAIA, M. (2019d). Extending Janzen’s

1096 hypothesis to temperate regions: a test using subterranean ecosystems. *Functional Ecology*

1097 **33**, 1638–1650.

1098 MARTÍNEZ, A., DI CESARE, A., MARI-MENA, N., GARCÍA-GÓMEZ, G., GARCIA-HERRERO, A.,
1099 CORNO, G., FONTANETO, D. & ECKERT, E.M. (2020). Tossed ‘good luck’ coins as vectors
1100 for anthropogenic pollution into aquatic environment. *Environmental Pollution* **259**,
1101 113800.

1102 MARTÍNEZ, A., KVINDEBJERG, K., ILIFFE, T.M. & WORSAAE, K. (2017). Evolution of cave
1103 suspension feeding in Protodrilidae (Annelida). *Zoologica Scripta* **46**, 214–226.

1104 MCGILL, B.J., CHASE, J.M., HORTAL, J., OVERCAST, I., ROMINGER, A.J., ROSINDELL, J., BORGES,
1105 P.A. V, EMERSON, B.C., ETIENNE, R.S., HICKERSON, M.J., MAHLER, D.L., MASSOL, F.,
1106 MCGAUGHRAN, A., NEVES, P., PARENT, C., *ET AL.* (2019). Unifying macroecology and
1107 macroevolution to answer fundamental questions about biodiversity. *Global Ecology and*
1108 *Biogeography* **28**, 1925–1936.

1109 MEDELLIN, R.A., WIEDERHOLT, R. & LOPEZ-HOFFMAN, L. (2017). Conservation relevance of bat
1110 caves for biodiversity and ecosystem services. *Biological Conservation* **211**, 45–50.

1111 MERMILLOD-BLONDIN, F. (2011). The functional significance of bioturbation and biodeposition
1112 on biogeochemical processes at the water–sediment interface in freshwater and marine
1113 ecosystems. *Journal of the North American Benthological Society* **30**, 770–778.

1114 MICHEL, G., MALARD, F., DEHARVENG, L., DI LORENZO, T., SKET, B. & DE BROYER, C. (2009).
1115 Reserve selection for conserving groundwater biodiversity. *Freshwater Biology* **54**, 861–
1116 876.

1117 MOKANY, K., HARWOOD, T.D., HALSE, S.A. & FERRIER, S. (2019). Riddles in the dark:
1118 Assessing diversity patterns for cryptic subterranean fauna of the Pilbara. *Diversity and*
1119 *Distributions* **25**, 240–254.

1120 MOLDOVAN, O.T., BERCEA, S., NĂSTASE-BUCUR, R., CONSTANTIN, S., KENESZ, M., MIREA, I.C.,
 1121 PETCULESCU, A., ROBU, M. & ARGHIR, R.A. (2020). Management of water bodies in show
 1122 caves – A microbial approach. *Tourism Management* **78**, 104037.

1123
 1124
 1125 MORIMURA, S., ZENG, X., NOBORU, N. & HOSONO, T. (2020). Changes to the microbial
 1126 communities within groundwater in response to a large crustal earthquake in Kumamoto,
 1127 southern Japan. *Journal of Hydrology* **581**, 124341.

1128 MORVAN, C., MALARD, F., PARADIS, E., LEFÉBURE, T., KONECNY-DUPRÉ, L. & DOUADY, C.J.
 1129 (2013). Timetree of aselloidea reveals species diversification dynamics in groundwater.
 1130 *Systematic Biology* **62**, 512–522.

1131 NIEMILLER, M.L., FITZPATRICK, B.M. & MILLER, B.T. (2008). Recent divergence with gene flow
 1132 in Tennessee cave salamanders (Plethodontidae: Gyrinophilus) inferred from gene
 1133 genealogies. *Molecular Ecology* **17**, 2258–2275.

1134 NIEMILLER, M.L., FITZPATRICK, B.M., SHAH, P., SCHMITZ, L. & NEAR, T.J. (2013). Evidence for
 1135 repeated loss of selective constraint in rhodopsin of amblyopsid cavefishes (Teleostei:
 1136 Amblyopsidae). *Evolution* **67**, 732–748.

1137 NIEMILLER, M.L., PORTER, M.L., KEANY, J., GILBERT, H., FONG, D.W., CULVER, D.C., HOBSON,
 1138 C.S., KENDALL, K.D., DAVIS, M.A. & TAYLOR, S.J. (2018). Evaluation of eDNA for
 1139 groundwater invertebrate detection and monitoring: a case study with endangered
 1140 *Stygobromus* (Amphipoda: Crangonyctidae). *Conservation Genetics Resources* **10**, 247–
 1141 257.

1142 NIEMILLER, M.L. & ZIGLER, K.S. (2013). Patterns of cave biodiversity and endemism in the

1143 Appalachians and Interior Plateau of Tennessee, USA. *PLoS ONE* **8**, e64177.

1144 NJUNJIĆ, I., PERRARD, A., HENDRIKS, K., SCHILTHUIZEN, M., PERREAU, M., MERCKX, V.,

1145 BAYLAC, M. & DEHARVENG, L. (2018). Comprehensive evolutionary analysis of the

1146 *Anthroherpon* radiation (Coleoptera, Leiodidae, Leptodirini). *PLoS ONE* **13**, e0198367.

1147 NORTHUP, D.E., MELIM, L.A., SPILDE, M.N., HATHAWAY, J.J.M., GARCIA, M.G., MOYA, M.,

1148 STONE, F.D., BOSTON, P.J., DAPKEVICIUS, M.L.N.E. & RIQUELME, C. (2011). Lava cave

1149 microbial communities within mats and secondary mineral deposits: implications for life

1150 detection on other planets. *Astrobiology* **11**, 601–618.

1151 ORTIZ, M., LEGATZKI, A., NEILSON, J.W., FRYSLIE, B., NELSON, W.M., WING, R.A., SODERLUND,

1152 C.A., PRYOR, B.M. & MAIER, R.M. (2014). Making a living while starving in the dark:

1153 metagenomic insights into the energy dynamics of a carbonate cave. *ISME Journal* **8**, 478–

1154 491.

1155 OZTURKOGLU-BUDAK, S., GURSOY, A., AYKAS, D.P., KOÇAK, C., DÖNMEZ, S., DE VRIES, R.P. &

1156 BRON, P.A. (2016). Volatile compound profiling of Turkish Divle Cave cheese during

1157 production and ripening. *Journal of Dairy Science* **99**, 5120–5131.

1158 PACIOGLU, O., IANOVICI, N., FILIMON, M., SINITEAN, A., IACOB, G., BARABAS, H., PAHOMI, A.,

1159 ACS, A., MUNTEAN, H. & PÂRVULESCU, L. (2019). The multifaceted effects induced by

1160 floods on the macroinvertebrate communities inhabiting a sinking cave stream.

1161 *International Journal of Speleology* **48**, 167–177.

1162 PATIÑO, J., WHITTAKER, R.J., BORGES, P.A. V, FERNÁNDEZ-PALACIOS, J.M., AH-PENG, C.,

1163 ARAÚJO, M.B., ÁVILA, S.P., CARDOSO, P., CORNUAULT, J., DE BOER, E.J., DE NASCIMENTO,

1164 L., GIL, A., GONZÁLEZ-CASTRO, A., GRUNER, D.S., HELENO, R., *ET AL.* (2017). A roadmap

1165 for island biology: 50 fundamental questions after 50 years of the Theory of Island

1166 Biogeography. *Journal of Biogeography* **44**, 963–983.

1167 PECK, S.B. & FINSTON, T.L. (1993). Galapagos islands troglobites: the questions of tropical
 1168 troglobites, parapatric distributions with eyed-sister-species, and their origin by parapatric
 1169 speciation. *Memoires de Biospeologie* **20**, 19–37.

1170 PÉREZ-MORENO, J.L., ILIFFE, T.M. & BRACKEN-GRISSOM, H.D. (2016). Life in the Underworld:
 1171 Anchialine cave biology in the era of speleogenomics. *International Journal of Speleology*
 1172 **49**, 149–170.

1173 PLAVÉN-SIGRAY, P., MATHESON, G.J., SCHIFFLER, B.C. & THOMPSON, W.H. (2017). The
 1174 readability of scientific texts is decreasing over time. *eLife* **6**, e27725.

1175 PONS, J., JURADO-RIVERA, J.A., JAUME, D., VONK, R., BAUZÀ-RIBOT, M.M. & JUAN, C. (2019).
 1176 The age and diversification of metacrangonyctid subterranean amphipod crustaceans
 1177 revisited. *Molecular Phylogenetics and Evolution* **140**, 10659.

1178 POPA, R., SMITH, A.R., POPA, R., BOONE, J. & FISK, M. (2011). Olivine-respiring bacteria
 1179 isolated from the rock-ice interface in a lava-tube cave, a mars analog environment.
 1180 *Astrobiology* **12**, 9–18.

1181 PORTER, M.L. & SUMNER-ROONEY, L. (2018). Evolution in the dark: unifying our understanding
 1182 of eye loss. *Integrative and Comparative Biology* **58**, 367–371.

1183 POULSON, T.L. & WHITE, W.B. (1969). The cave environment. *Science* **165**, 971–981.

1184 PRENDINI, L., FRANCKE, O.F. & VIGNOLI, V. (2010). Troglomorphism, trichobothriotaxy and
 1185 typhlochactid phylogeny (Scorpiones, Chactioidea): more evidence that troglobitism is not
 1186 an evolutionary dead-end. *Cladistics* **26**, 117–142.

1187 PROTAS, M. & JEFFERY, W.R. (2012). Evolution and development in cave animals: from fish to
 1188 crustaceans. *Wiley interdisciplinary reviews. Developmental biology* **1**, 823–845.

1189 PYŠEK, P., HULME, P.E., SIMBERLOFF, D., BACHER, S., BLACKBURN, T.M., CARLTON, J.T.,
 1190 DAWSON, W., ESSL, F., FOXCROFT, L.C., GENOVESI, P., JESCHKE, J.M., KÜHN, I., LIEBHOLD,
 1191 A.M., MANDRAK, N.E., MEYERSON, L.A., *ET AL.* (2020). Scientists' warning on invasive
 1192 alien species. *Biological Reviews*, 10.1111/brv.12627
 1193 RABELO, L.M., SOUZA-SILVA, M. & FERREIRA, R.L. (2018). Priority caves for biodiversity
 1194 conservation in a key karst area of Brazil: comparing the applicability of cave conservation
 1195 indices. *Biodiversity and Conservation* **27**, 2097–2129.
 1196 RAHMSTORF, S. & COUMOU, D. (2011). Increase of extreme events in a warming world.
 1197 *Proceedings of the National Academy of Sciences* **108**, 17905–17909.
 1198 REBOLEIRA, A.S.P.S., ABRANTES, N., OROMÍ, P. & GONÇALVES, F. (2013). Acute toxicity of
 1199 copper sulfate and potassium dichromate on stygobiont *Proasellus*: general aspects of
 1200 groundwater ecotoxicology and future perspectives. *Water, Air, & Soil Pollution* **224**, 1550.
 1201 RIDDLE, M.R., ASPIRAS, A.C., GAUDENZ, K., PEUB, R., SUNG, J.Y., MARTINEAU, B., PEAVEY, M.,
 1202 BOX, A.C., TABIN, J.A., MCGAUGH, S., BOROWSKY, R., TABIN, C.J. & ROHNER, N. (2018).
 1203 Insulin resistance in cavefish as an adaptation to a nutrient-limited environment. *Nature*
 1204 **555**, 647.
 1205 RIESCH, R., REZNICK, D.N., PLATH, M. & SCHLUPP, I. (2016). Sex-specific local life-history
 1206 adaptation in surface- and cave-dwelling Atlantic mollies (*Poecilia mexicana*). *Scientific*
 1207 *Reports* **6**, 22968.
 1208 RIPPLE, W.J., WOLF, C., NEWSOME, T.M., BARNARD, P. & MOOMAW, W.R. (2019). World
 1209 scientists' warning of a climate emergency. *BioScience* **70**, 8–12.
 1210 RIPPLE, W.J., WOLF, C., NEWSOME, T.M., GALETTI, M., ALAMGIR, M., CRIST, E., MAHMOUD,
 1211 M.I., LAURANCE, W.F. & 15364 SCIENTIST SIGNATORIES FROM 184 COUNTRIES (2017).

1212 World scientists' warning to humanity: a second notice. *BioScience* **67**, 1026–1028.

1213 SAMWAYS, M.J., BARTON, P.S., BIRKHOFFER, K., CHICHORRO, F., DEACON, C., FARTMANN, T.,
1214 FUKUSHIMA, C.S., GAIGHER, R., HABEL, J.C., HALLMANN, C.A., HILL, M.J., HOCHKIRCH, A.,
1215 KAILA, L., KWAK, M.L., MAES, D., *ET AL.* (2020). Solutions for humanity on how to
1216 conserve insects. *Biological Conservation* **242**, 108427.

1217 SÁNCHEZ-FERNÁNDEZ, D., RIZZO, V., BOURDEAU, C., CIESLAK, A., COMAS, J., FAILLE, A.,
1218 FRESNEDA, J., LLEOPART, E., MILLÁN, A., MONTES, A., PALLARES, S. & RIBERA, I. (2018).
1219 The deep subterranean environment as a potential model system in ecological,
1220 biogeographical and evolutionary research. *Subterranean Biology* **25**, 1–7.

1221 SARBU, S.M., KANE, T.C. & KINKLE, B.K. (1996). A chemoautotrophically based cave
1222 ecosystem. *Science* **272**, 1953–1955.

1223 SCHEFFER, M., CARPENTER, S., FOLEY, J.A., FOLKE, C. & WALKER, B. (2001). Catastrophic
1224 shifts in ecosystems. *Nature* **413**, 591–596.

1225 SEDDON, A.W.R., MACKAY, A.W., BAKER, A.G., BIRKS, H.J.B., BREMAN, E., BUCK, C.E., ELLIS,
1226 E.C., FROYD, C.A., GILL, J.L., GILLSON, L., JOHNSON, E.A., JONES, V.J., JUGGINS, S.,
1227 MACIAS-FAURIA, M., MILLS, K., *ET AL.* (2014). Looking forward through the past:
1228 identification of 50 priority research questions in palaeoecology. *Journal of Ecology* **102**,
1229 256–267.

1230 SIMIČEVIĆ, V. (2017). Poachers threaten Balkans underground biodiversity. *Science* **358**, 1116
1231 LP – 1117.

1232 SIMON, V., ELLEBOODE, R., MAHÉ, K., LEGENDRE, L., ORNELAS-GARCIA, P., ESPINASA, L. &
1233 RÉTAUX, S. (2017). Comparing growth in surface and cave morphs of the species *Astyanax*
1234 *mexicanus*: insights from scales. *EvoDevo* **8**, 23.

1235 SMITH, V.H. (2007). Microbial diversity-productivity relationships in aquatic ecosystems. *FEMS*
1236 *Microbiology Ecology* **62**, 181–186.

1237 STERANRS, S. (1992). *The Evolution of Life Histories*. Oxford University Press, New York.

1238 STERN, D.B., BREINHOLT, J., PEDRAZA-LARA, C., LÓPEZ-MEJÍA, M., OWEN, C.L., BRACKEN-
1239 GRISSOM, H., FETZNER JR., J.W. & CRANDALL, K.A. (2017). Phylogenetic evidence from
1240 freshwater crayfishes that cave adaptation is not an evolutionary dead-end. *Evolution* **71**,
1241 2522–2532.

1242 STRONA, G., FATTORINI, S., FIASCA, B., DI LORENZO, T., DI CICCIO, M., LORENZETTI, W.,
1243 BOCCACCI, F. & GALASSI, D.M.P. (2019). Aqualife software: a new tool for a standardized
1244 ecological assessment of groundwater dependent ecosystems. *Water* **11**, 2574.

1245 SUTHERLAND, W.J., DIAS, M.P., DICKS, L. V, DORAN, H., ENTWISTLE, A.C., FLEISHMAN, E.,
1246 GIBBONS, D.W., HAILS, R., HUGHES, A.C., HUGHES, J., KELMAN, R., LE ROUX, X.,
1247 LEANSTEY, B., LICKORISH, F.A., MAGGS, L., *ET AL.* (2020). A horizon scan of emerging
1248 global biological conservation issues for 2020. *Trends in Ecology & Evolution* **35**, 81–90.

1249 SUTHERLAND, W.J., FLEISHMAN, E., MASCIA, M.B., PRETTY, J. & RUDD, M.A. (2011). Methods
1250 for collaboratively identifying research priorities and emerging issues in science and policy.
1251 *Methods in Ecology and Evolution* **2**, 238–247.

1252 SUTHERLAND, W.J., FRECKLETON, R.P., GODFRAY, H.C.J., BEISSINGER, S.R., BENTON, T.,
1253 CAMERON, D.D., CARMEL, Y., COOMES, D.A., COULSON, T., EMMERSON, M.C., HAILS, R.S.,
1254 HAYS, G.C., HODGSON, D.J., HUTCHINGS, M.J., JOHNSON, D., *ET AL.* (2013). Identification of
1255 100 fundamental ecological questions. *Journal of Ecology* **101**, 58–67.

1256 TANALGO, K.C., TABORA, J.A.G. & HUGHES, A.C. (2018). Bat cave vulnerability index (BCVI):
1257 a holistic rapid assessment tool to identify priorities for effective cave conservation in the

1258 tropics. *Ecological Indicators* **89**, 852–860.

1259 TIERNEY, S.M., LANGILLE, B., HUMPHREYS, W.F., AUSTIN, A.D. & COOPER, S.J.B. (2018).

1260 Massive parallel regression: a précis of genetic mechanisms for vision loss in diving beetles.

1261 *Integrative and Comparative Biology* **58**, 465–479.

1262 TOBIN, B.W., HUTCHINS, B.T. & SCHWARTZ, B.F. (2013). Spatial and temporal changes in

1263 invertebrate assemblage structure from the entrance to deep-cave zone of a temperate

1264 marble cave. *International Journal of Speleology* **42**, 203–214.

1265 TORRES-PAZ, J., HYACINTHE, C., PIERRE, C. & RÉTAUX, S. (2018). Towards an integrated

1266 approach to understand Mexican cavefish evolution. *Biology Letters* **14**, 20180101.

1267 TRONTELJ, P. (2019). Structure and genetics of cave populations. In *Cave Ecology* (eds T.

1268 MOLDOVAN, L. KOVÁČ & S. HALSE), pp. 269–296. Springer, Cham, Switzerland.

1269 TRONTELJ, P., BLEJEC, A. & FIŠER, C. (2012). Ecomorphological convergence of cave

1270 communities. *Evolution* **66**, 3852–3865.

1271 TRONTELJ, P., BORKO, Š. & DELIĆ, T. (2019). Testing the uniqueness of deep terrestrial life.

1272 *Scientific Reports* **9**, 15188.

1273 VENARSKY, M.P. & HUNTSMAN, B.M. (2018). Food webs in caves. In *Cave Ecology* (eds O.

1274 MOLDOVAN, L. KOVÁČ & S. HALSE), pp. 309–328. Springer, Cham, Switzerland.

1275 VENARSKY, M.P., HURYN, A.D. & BENSTEAD, J.P. (2012). Re-examining extreme longevity of

1276 the cave crayfish *Orconectes australis* using new mark–recapture data: a lesson on the

1277 limitations of iterative size-at-age models. *Freshwater Biology* **57**, 1471–1481.

1278 VOITURON, Y., DE FRAIPONT, M., ISSARTEL, J., GUILLAUME, O. & CLOBERT, J. (2011). Extreme

1279 lifespan of the human fish (*Proteus anguinus*): A challenge for ageing mechanisms. *Biology*

1280 *Letters*, 7.

1281 VONK, R. & NIJMAN, V. (2006). Sex ratio and sexual selection in wormshrimps (Crustacea,
1282 Amphipoda, Ingolfiellidea). *Contributions to Zoology* **75**, 189–194.

1283 WAKELING, S., WILLETT, P., CREASER, C., FRY, J., PINFIELD, S. & SPEZI, V. (2016). Open-access
1284 mega-journals: a bibliometric profile. *PLoS ONE* **11**, e0165359.

1285 WALTHER, G.-R., POST, E., CONVEY, P., MENZEL, A., PARMESAN, C., BEEBEE, T.J.C.,
1286 FROMENTIN, J.-M., HOEGH-GULDBERG, O. & BAIRLEIN, F. (2002). Ecological responses to
1287 recent climate change. *Nature* **416**, 389–395.

1288 WILKENS, H. (2020). The role of selection in the evolution of blindness in cave fish. *Biological*
1289 *Journal of the Linnean Society* **130**, 421–432.

1290 WILKENS, H. & STRECKER, U. (2017). *Evolution in the dark: Darwin's loss without selection*.
1291 Springer, Cham, Switzerland.

1292 WYNNE, J.J., BERNARD, E.C., HOWARTH, F.G., SOMMER, S., SOTO-ADAMES, F.N., TAITI, S.,
1293 MOCKFORD, E.L., HORROCKS, M., PAKARATI, L. & PAKARATI-HOTUS, V. (2014).
1294 Disturbance relicts in a rapidly changing world: the Rapa Nui (Easter Island) factor.
1295 *BioScience* **64**, 711–718.

1296 WYNNE, J.J., HOWARTH, F.G., SOMMER, S. & DICKSON, B.G. (2019). Fifty years of cave
1297 arthropod sampling: techniques and best practices. *International Journal of Speleology* **48**,
1298 33–48.

1299 WYNNE, J.J., SOMMER, S., HOWARTH, F.G., DICKSON, B.G. & VOYLES, K.D. (2018). Capturing
1300 arthropod diversity in complex cave systems. *Diversity and Distributions* **24**, 1478–1491.

1301 YOSHIZAWA, M., GORIČKI, Š., SOARES, D. & JEFFERY, W.R. (2010). Evolution of a behavioral
1302 shift mediated by superficial neuromasts helps cavefish find food in darkness. *Current*
1303 *Biology* **20**, 1631–1636.

- 1304 YOSHIZAWA, M., SETTLE, A., HERMOSURA, M.C., TUTTLE, L.J., CETRARO, N., PASSOW, C.N. &
 1305 MCGAUGH, S.E. (2018). The evolution of a series of behavioral traits is associated with
 1306 autism-risk genes in cavefish. *BMC Evolutionary Biology* **18**, 89.
- 1307 ZAGMAJSTER, M., CULVER, D.C. & SKET, B. (2008). Species richness patterns of obligate
 1308 subterranean beetles (Insecta: Coleoptera) in a global biodiversity hotspot – effect of scale
 1309 and sampling intensity. *Diversity and Distributions* **14**, 95–105.
- 1310 ZAGMAJSTER, M., EME, D., FIŠER, C., GALASSI, D., MARMONIER, P., STOCH, F., CORNU, J.F. &
 1311 MALARD, F. (2014). Geographic variation in range size and beta diversity of groundwater
 1312 crustaceans: Insights from habitats with low thermal seasonality. *Global Ecology and*
 1313 *Biogeography* **23**, 1135–1145.
- 1314 ZAGMAJSTER, M., MALARD, F., EME, D. & CULVER, D.C. (2019). Subterranean biodiversity
 1315 patterns from global to regional scales. In *Cave Ecology* (eds T. MOLDOVAN, L. KOVÁČ &
 1316 S. HALSE) pp. 195–227. Springer, Cham, Switzerland.
- 1317 ZAKŠEK, V., DELIĆ, T., FIŠER, C., JALŽIĆ, B. & TRONTELJ, P. (2019). Emergence of sympatry in a
 1318 radiation of subterranean amphipods. *Journal of Biogeography* **46**, 657–669.
- 1319 ZHANG, J., KAPLI, P., PAVLIDIS, P. & STAMATAKIS, A. (2013). A general species delimitation
 1320 method with applications to phylogenetic placements. *Bioinformatics* **29**, 2869–2876.

1321

1322 **XIII. SUPPORTING INFORMATION**

1323 Additional supporting information may be found online in the Supporting Information section at
 1324 the end of the article.

1325 **Appendix S1.** Questions from List #2 (i.e. 120 questions selected from List #1 during Survey#1)
 1326 and List #3 (i.e. 25 additional questions suggested by Survey #2 participants) ranked based on

1327 the percentage of ‘major importance’ votes.

1328

1329 Table 1. Subject areas, general topics addressed, panel member composition (*= panel
 1330 coordinator; °= postdoc or early career researcher), and number of questions included in the top-
 1331 50 list out of the total retained in List #1. Panel members are listed alphabetically by surname.
 1332

Subject area	General topics	Panel members	Number of questions
Adaptation	Morphological, physiological and behavioural adaptations to the subterranean environment	Žiga Fišer°, Daniel W. Fong, Tanja Pipan*, William R. Jeffery, Jure Jugovic	10 out of 43
Origin and evolution	Cave ontology and past climate change, migration–speciation–extinction dynamics, and speciation and diversification	Steven J.B. Cooper*, Matthew Niemiller, Alejandro Martínez°, Meredith Protas	11 out of 36
Community ecology	Population dynamics, community assembly, biotic interaction, trophic webs, and energy flows	Rodrigo L. Ferreira*, Cene Fišer, Thais G. Pellegrini°, Michael Venarsky°	4 out of 32
Macroecology and biogeography	Global diversity patterns (taxonomic, phylogenetic, functional), biogeography theory, and diversity drivers	Maria E. Bichuette, David Eme°, Florian Malard*, Maja Zgamažster°	6 out of 32
Conservation biology	Climate change, habitat loss, invasive species, conservation and management policies, and show-cave-related issues	Isabel R. Amorim°, Paulo A. V. Borges*, Louis Deharveng, J. Judson Wynne, Ana Sofia P. S. Reboleira	12 out of 37
Microbiology and applied topics	Microbial communities, industrial and pharmaceutical potential, epidemics, and exobiology	Naowarat Cheeptham, Thomas M. Lilley*, Melissa B. Meierhofer°, Diana E. Northup	7 out of 31
Other topics	Any topic falling outside the scope of the six core subject areas	David C. Culver*, Christian Griebler, Johanna Kowalko, Raoul Manenti°	n/a (merged within the other subject areas)

1333

1334 Table 2. Glossary of terms.

1335

Term	General definition
Cave	A human-accessible subterranean space, either a single chamber or series of chambers, formed within different substrata (Curl, 1964). Note that a cave is just one among the wide variety of subterranean habitats (see definition below).
Exaptation	A trait shaped by selection or neutral evolution co-opted for a new function (Gould & Vrba, 1982).
Speleogenetic process	The process of water dissolving surrounding rock, gradually forming passages that evolve into cave systems (Audra & Palmer, 2011).
Subterranean habitat(s) / ecosystem(s)	The breadth of underground voids of different sizes, either dry or filled with water, sharing two main ecological features: the absence of sunlight and buffered climatic conditions. Examples of subterranean habitats include caves, groundwater, anchialine systems, artificially excavated underground voids, shallow subterranean habitats, as well as deep maze of fissures and pore spaces with size prohibiting human entry (Culver & Pipan, 2019).

1336

FIGURE LEGENDS

Fig. 1. Survey workflow, summary statistics of survey participants, and the breakdown by subject area of the 50 highest priority research questions.

Fig. 2. The relationship between median range size (maximum linear extent) per latitudinal band and latitude for 147 European groundwater species of Niphargidae (Amphipoda) and Aselloidea (Isopoda) delimited using morphology (A) and a molecular species delimitation method (B). Molecular delimitation was performed by a Bayesian implementation of the Poisson tree processes (Zhang *et al.*, 2013) approach based on molecular phylogenies inferred from 2883 cytochrome *c* oxidase subunit I sequences. Black horizontal bars, dots, and boxes show the median, average, and interquartile range, respectively, for 0.9° latitudinal bands. The maximum length of each whisker is up to 1.5 times the interquartile range. Trend lines (with 95% confidence intervals) represent the fit of a gamma generalized linear model to the averages of latitudinal bands and its quadratic (A) and cubic (B) term. Data re-analysed from Eme *et al.* (2018).